

RBP-L siRNA (m): sc-152762

BACKGROUND

Recombination signal binding protein for immunoglobulin κ J region-like (RBP-L), whose alternative names include recombining binding protein suppressor of hairless-like protein, SUH, RBPJL, SUHL and RBPSUHL, is a 516 amino acid transcription factor belonging to the suppressor of hairless [Su(H)] family. RBP-L is closely related to RBP-J κ (recombination signal binding protein J κ), a transcription factor involved in the Notch signaling pathway. RBP-L binds DNA sequences nearly identical to that of RBP-J κ , however, RBP-L does not interact with Notch receptors. RBP-L localizes to the nucleus and is known to interact weakly with EBNA2 (Epstein-Barr virus nuclear antigen-2) to activate transcription. RBP-L contains one IPT/TIG domain and shows almost exclusive expression in lung. The gene encoding RBP-L maps to human chromosome 20q13.12.

REFERENCES

1. Amakawa, R., Jing, W., Ozawa, K., Matsunami, N., Hamaguchi, Y., Matsuda, F., Kawaichi, M. and Honjo, T. 1993. Human J κ recombination signal binding protein gene (IGKJRB): comparison with its mouse homologue. *Genomics* 17: 306-315.
2. Waltzer, L., Logeat, F., Brou, C., Israel, A., Sergeant, A. and Manet, E. 1994. The human J κ recombination signal sequence binding protein (RBP-J κ) targets the Epstein-Barr virus EBNA2 protein to its DNA responsive elements. *EMBO J.* 13: 5633-5638.
3. Minoguchi, S., Taniguchi, Y., Kato, H., Okazaki, T., Strobl, L.J., Zimmer-Strobl, U., Bornkamm, G.W. and Honjo, T. 1997. RBP-L, a transcription factor related to RBP-J κ . *Mol. Cell. Biol.* 17: 2679-2687.
4. Tani, S., Taniwaki, M., Taniguchi, Y., Minoguchi, S., Kuroda, K., Han, H., Aoki, T., Miyatake, S., Hashimoto, N. and Honjo, T. 1999. Chromosomal mapping of two RBP-J-related genes: Kyo-T and RBP-L. *J. Hum. Genet.* 44: 73-75.
5. Beres, T.M., Masui, T., Swift, G.H., Shi, L., Henke, R.M. and MacDonald, R.J. 2006. PTF1 is an organ-specific and Notch-independent basic helix-loop-helix complex containing the mammalian suppressor of hairless (RBP-J) or its paralogue, RBP-L. *Mol. Cell. Biol.* 26: 117-130.

CHROMOSOMAL LOCATION

Genetic locus: Rbpjl (mouse) mapping to 2 H3.

PRODUCT

RBP-L siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RBP-L shRNA Plasmid (m): sc-152762-SH and RBP-L shRNA (m) Lentiviral Particles: sc-152762-V as alternate gene silencing products.

For independent verification of RBP-L (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152762A, sc-152762B and sc-152762C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RBP-L siRNA (m) is recommended for the inhibition of RBP-L expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RBP-L gene expression knockdown using RT-PCR Primer: RBP-L (m)-PR: sc-152762-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.