



Ribosomal Protein L38 siRNA (m): sc-152921

BACKGROUND

Ribosomes, the organelles that catalyze protein synthesis, are composed of a small subunit (40S) and a large subunit (60S) that consist of over 80 distinct ribosomal proteins. Mammalian ribosomal proteins are encoded by multigene families that contain processed pseudogenes and one functional intron-containing gene within their coding regions. Ribosomal Protein L38 (RPL38) is a 70 amino acid protein that localizes to the cytoplasm and functions as a component of the 60S subunit where it likely plays a role in protein translation. RPL38 belongs to the Ribosomal Protein L38E family of ribosomal proteins and may be regulated by GABP. Like most ribosomal proteins, Ribosomal Protein L38 exists as multiple processed pseudogenes that are scattered throughout the genome.

REFERENCES

1. Ou, J.H., Yen, T.S., Wang, Y.F., Kam, W.K. and Rutter, W.J. 1987. Cloning and characterization of a human ribosomal protein gene with enhanced expression in fetal and neoplastic cells. *Nucleic Acids Res.* 15: 8919-8934.
2. Reddy, T.R., Suhasini, M., Rappaport, J., Looney, D.J., Kraus, G. and Wong-Staal, F. 1995. Molecular cloning and characterization of a TAR-binding nuclear factor from T cells. *AIDS Res. Hum. Retroviruses* 11: 663-669.
3. Wool, I.G., Chan, Y.L. and Glück, A. 1995. Structure and evolution of mammalian ribosomal proteins. *Biochem. Cell Biol.* 73: 933-947.
4. Kenmochi, N., Kawaguchi, T., Rozen, S., Davis, E., Goodman, N., Hudson, T.J., Tanaka, T. and Page, D.C. 1998. A map of 75 human ribosomal protein genes. *Genome Res.* 8: 509-523.
5. Duga, S., Asselta, R., Malcovati, M., Tenchini, M.L., Ronchi, S. and Simonini, T. 2000. The intron-containing L3 ribosomal protein gene (RPL3): sequence analysis and identification of U43 and of two novel intronic small nucleolar RNAs. *Biochim. Biophys. Acta* 1490: 225-236.
6. Odintsova, T.I., Müller, E.C., Ivanov, A.V., Egorov, T.A., Bienert, R., Vladimirov, S.N., Kostka, S., Otto, A., Wittmann-Liebold, B. and Karpova, G.G. 2003. Characterization and analysis of posttranslational modifications of the human large cytoplasmic ribosomal subunit proteins by mass spectrometry and Edman sequencing. *J. Protein Chem.* 22: 249-258.
7. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 604163. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Rpl38 (mouse) mapping to 11 E2.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Ribosomal Protein L38 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Ribosomal Protein L38 shRNA Plasmid (m): sc-152921-SH and Ribosomal Protein L38 shRNA (m) Lentiviral Particles: sc-152921-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Ribosomal Protein L38 siRNA (m) is recommended for the inhibition of Ribosomal Protein L38 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.