

RMND5A siRNA (m): sc-152985

BACKGROUND

RMND5A (required for meiotic nuclear division 5 homolog A), also known as CTLH (C-terminal to the Lissencephaly type-1-like homology), RMD5 or p44CTLH, is a 391 amino acid protein containing one CTLH domain and one LisH domain. Existing as two alternatively spliced isoforms, RMND5A plays an essential role in the degradation of fructose-1,6-bisphosphatase (FR) in yeast and is well conserved in mammalian cells. RMND5A is encoded by a gene located on human chromosome 2, which houses over 1,400 genes and comprises nearly 8% of the human genome. Harlequin ichthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene, while the lipid metabolic disorder sitosterolemia is associated with defects in the ABCG5 and ABCG8 genes. Additionally, an extremely rare recessive genetic disorder, Alström syndrome, is caused by mutations in the ALMS1 gene, which maps to chromosome 2.

REFERENCES

1. Ijdo, J.W., Baldini, A., Ward, D.C., Reeders, S.T. and Wells, R.A. 1991. Origin of human chromosome 2: an ancestral telomere-telomere fusion. *Proc. Natl. Acad. Sci. USA* 88: 9051-9055.
2. Avarello, R., Pedicini, A., Caiulo, A., Zuffardi, O. and Fraccaro, M. 1992. Evidence for an ancestral alphoid domain on the long arm of human chromosome 2. *Hum. Genet.* 89: 247-249.
3. Umeda, M., Nishitani, H. and Nishimoto, T. 2003. A novel nuclear protein, Twa1, and Muskelein comprise a complex with RanBPM. *Gene* 303: 47-54.
4. Regelman, J., Schüle, T., Josupeit, F.S., Horak, J., Rose, M., Entian, K.D., Thumm, M. and Wolf, D.H. 2003. Catabolite degradation of fructose-1,6-bisphosphatase in the yeast *Saccharomyces cerevisiae*: a genome-wide screen identifies eight novel GID genes and indicates the existence of two degradation pathways. *Mol. Biol. Cell* 14: 1652-1663.
5. Kobayashi, N., Yang, J., Ueda, A., Suzuki, T., Tomaru, K., Takeno, M., Okuda, K. and Ishigatsubo, Y. 2007. RanBPM, Muskelein, p48EMLP, p44CTLH, and the armadillo-repeat proteins ARMC8 α and ARMC8 β are components of the CTLH complex. *Gene* 396: 236-247.
6. Suzuki, T., Ueda, A., Kobayashi, N., Yang, J., Tomaru, K., Yamamoto, M., Takeno, M. and Ishigatsubo, Y. 2008. Proteasome-dependent degradation of α -catenin is regulated by interaction with ARMC8 α . *Biochem. J.* 411: 581-591.
7. Tomaru, K., Ueda, A., Suzuki, T., Kobayashi, N., Yang, J., Yamamoto, M., Takeno, M., Kaneko, T. and Ishigatsubo, Y. 2010. Armadillo repeat containing 8 α binds to HRS and promotes HRS interaction with ubiquitinated proteins. *Open Biochem. J.* 4: 1-8.

CHROMOSOMAL LOCATION

Genetic locus: Rmnd5a (mouse) mapping to 6 C1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

RMND5A siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RMND5A shRNA Plasmid (m): sc-152985-SH and RMND5A shRNA (m) Lentiviral Particles: sc-152985-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RMND5A siRNA (m) is recommended for the inhibition of RMND5A expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RMND5A gene expression knockdown using RT-PCR Primer: RMND5A (m)-PR: sc-152985-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.