

RNF11 siRNA (m): sc-153001

BACKGROUND

The RING-type zinc-finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in the ubiquitination pathway of protein degradation. RNF11 (RING finger protein 11), also known as CGI-123, is a 154 amino acid protein that contains one RING-type zinc-finger through which it interacts with a variety of proteins. Specifically, RNF11 associates with NEDD4 (an E3 ubiquitin-protein ligase) and EGFR (epidermal growth factor receptor) and may regulate the function of these proteins. Via its ability to control the activity of growth-associated proteins, such as EGFR, RNF11 is associated with the pathogenesis of endocrine neoplasia, Parkinson's disease and breast cancer.

REFERENCES

1. Seki, N., et al. 1999. Cloning and expression profile of mouse and human genes, Rnf11/RNF11, encoding a novel RING-H2 finger protein. *Biochim. Biophys. Acta* 1489: 421-427.
2. Jolliffe, C.N., et al. 2000. Identification of multiple proteins expressed in murine embryos as binding partners for the WW domains of the ubiquitin-protein ligase NEDD4. *Biochem. J.* 351: 557-565.
3. Kitching, R., et al. 2003. The RING-H2 protein RNF11 is differentially expressed in breast tumours and interacts with HECT-type E3 ligases. *Biochim. Biophys. Acta* 1639: 104-112.
4. Subramaniam, V., et al. 2003. The RING-H2 protein RNF11 is overexpressed in breast cancer and is a target of Smurf2 E3 ligase. *Br. J. Cancer* 89: 1538-1544.
5. Li, H. and Seth, A. 2004. An RNF11: Smurf2 complex mediates ubiquitination of the AMSH protein. *Oncogene* 23: 1801-1808.
6. Connor, M.K., et al. 2005. Molecular characterization of RING finger protein 11. *Mol. Cancer Res.* 3: 453-461.
7. Burger, A., et al. 2006. Novel RING-E3 ubiquitin ligases in breast cancer. *Neoplasia* 8: 689-695.
8. Anderson, L.R., et al. 2007. PARK10 candidate RNF11 is expressed by vulnerable neurons and localizes to Lewy bodies in Parkinson disease brain. *J. Neuropathol. Exp. Neurol.* 66: 955-964.

CHROMOSOMAL LOCATION

Genetic locus: Rnf11 (mouse) mapping to 4 C7.

PRODUCT

RNF11 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RNF11 shRNA Plasmid (m): sc-153001-SH and RNF11 shRNA (m) Lentiviral Particles: sc-153001-V as alternate gene silencing products.

For independent verification of RNF11 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153001A, sc-153001B and sc-153001C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCL, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RNF11 siRNA (m) is recommended for the inhibition of RNF11 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

RNF11 (4G7): sc-517151 is recommended as a control antibody for monitoring of RNF11 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RNF11 gene expression knockdown using RT-PCR Primer: RNF11 (m)-PR: sc-153001-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.