



RNF32 siRNA (m): sc-153047

BACKGROUND

The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in the ubiquitination pathway of protein degradation. RNF32 (RING finger protein 32), also known as HSD15 or FKSG33, is a 362 amino acid cytoplasmic protein that contains one IQ domain and two RING-type zinc fingers. Highly expressed in testis with lower expression levels in ovary tissue, RNF32 is thought to play a role in spermatogenesis, specifically contributing to the growth and maturation of round spermatids. Six isoforms of RNF32 exist due to alternative splicing events.

REFERENCES

1. Borden, K.L. and Freemont, P.S. 1996. The RING finger domain: a recent example of a sequence-structure family. *Curr. Opin. Struct. Biol.* 6: 395-401.
2. Lorick, K.L., Jensen, J.P., Fang, S., Ong, A.M., Hatakeyama, S. and Weissman, A.M. 1999. RING fingers mediate ubiquitin-conjugating enzyme (E2)-dependent ubiquitination. *Proc. Natl. Acad. Sci. USA* 96: 11364-11369.
3. van Baren, M.J., van der Linde, H.C., Breedveld, G.J., Baarends, W.M., Rizzu, P., de Graaff, E., Oostra, B.A. and Heutink, P. 2002. A double RING-H2 domain in RNF32, a gene expressed during sperm formation. *Biochem. Biophys. Res. Commun.* 292: 58-65.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 610241. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Colland, F., Jacq, X., Trouplin, V., Mougin, C., Groizeleau, C., Hamburger, A., Meil, A., Wojcik, J., Legrain, P. and Gauthier, J.M. 2004. Functional proteomics mapping of a human signaling pathway. *Genome Res.* 14: 1324-1332.
6. Penengo, L., Mapelli, M., Murachelli, A.G., Confalonieri, S., Magri, L., Musacchio, A., Di Fiore, P.P., Polo, S. and Schneider, T.R. 2006. Crystal structure of the ubiquitin binding domains of rabex-5 reveals two modes of interaction with ubiquitin. *Cell* 124: 1183-1195.

CHROMOSOMAL LOCATION

Genetic locus: Rnf32 (mouse) mapping to 5 B1.

PRODUCT

RNF32 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RNF32 shRNA Plasmid (m): sc-153047-SH and RNF32 shRNA (m) Lentiviral Particles: sc-153047-V as alternate gene silencing products.

For independent verification of RNF32 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153047A, sc-153047B and sc-153047C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RNF32 siRNA (m) is recommended for the inhibition of RNF32 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RNF32 gene expression knockdown using RT-PCR Primer: RNF32 (m)-PR: sc-153047-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.