

SMR3A siRNA (m): sc-153639

BACKGROUND

Salivary glands are accessory digestive structures that also are sources of systemically active immunoregulatory and anti-inflammatory factors. The salivary glands participate in a neuroendocrine axis that is regulated by the autonomic nervous system and contributes to whole body homeostasis. Submaxillary gland androgen-regulated proteins are male-specific proteins that have a prohormone structure and are processed by maturation enzymes, leading to the release of small peptides within the blood and saliva. These peptides have analgesic and anti-inflammatory properties, and seem to be important in mediating inflammation, suppressing pain and erectile dysfunction. Submaxillary gland androgen-regulated protein 3A (SMR3A) and 3B (SMR3B), also known as proline-rich protein 5 and proline-rich protein 3 respectively, are secreted into the saliva by the submaxillary gland and may have a hormonal effect.

REFERENCES

1. Rosinski-Chupin, I., Rougeot, C., Courty, Y. and Rougeon, F. 1993. Localization of mRNAs of two androgen-dependent proteins, SMR1 and SMR2, by *in situ* hybridization reveals sexual differences in acinar cells of rat submandibular gland. *J. Histochem. Cytochem.* 41: 1645-1649.
2. Tronik-Le Roux, D., Señorale-Pose, M. and Rougeon, F. 1994. Three novel SMR1-related cDNAs characterized in the submaxillary gland of mice show extensive evolutionary divergence in the protein coding region. *Gene* 142: 175-182.
3. Singer, M., Courty, Y. and Rougeon, F. 1995. Recent evolution of genes encoding the prohormone-like protein SMR1 in the rat submandibular gland. *DNA Cell Biol.* 14: 137-144.
4. Isemura, S. and Saitoh, E. 1997. Nucleotide sequence of gene PBI encoding a protein homologous to salivary proline-rich protein P-B. *J. Biochem.* 121: 1025-1030.
5. Señorale-Pose, M., Jacqueson, A., Rougeon, F. and Rosinski-Chupin, I. 1998. Acinar cells are target cells for androgens in mouse submandibular glands. *J. Histochem. Cytochem.* 46: 669-678.
6. Rougeot, C., Rosinski-Chupin, I., Mathison, R. and Rougeon, F. 2000. Rodent submandibular gland peptide hormones and other biologically active peptides. *Peptides* 21: 443-455.
7. User, H.M., Zelner, D.J., McKenna, K.E. and McVary, K.T. 2003. Microarray analysis and description of SMR1 gene in rat penis in a post-radical prostatectomy model of erectile dysfunction. *J. Urol.* 170: 298-301.
8. Morris, K.E., St Laurent, C.D., Hoeve, R.S., Forsythe, P., Suresh, M.R., Mathison, R.D. and Befus, A.D. 2009. Autonomic nervous system regulates secretion of anti-inflammatory prohormone SMR1 from rat salivary glands. *Am. J. Physiol., Cell Physiol.* 296: C514-C524.
9. Mathison, R.D., Davison, J.S., Befus, A.D. and Gingerich, D.A. 2010. Salivary gland derived peptides as a new class of anti-inflammatory agents: review of preclinical pharmacology of C-terminal peptides of SMR1 protein. *J. Inflamm.* 7: 49.

CHROMOSOMAL LOCATION

Genetic locus: *Smr3a* (mouse) mapping to 5 E1.

PRODUCT

SMR3A siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SMR3A shRNA Plasmid (m): sc-153639-SH and SMR3A shRNA (m) Lentiviral Particles: sc-153639-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SMR3A siRNA (m) is recommended for the inhibition of SMR3A expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SMR3A gene expression knockdown using RT-PCR Primer: SMR3A (m)-PR: sc-153639-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.