

SNX33 siRNA (m): sc-153678

BACKGROUND

SNX33 (sorting nexin-33), also known as SH3PX3, SH3PXD3C or SNX30, is a 574 amino acid protein that interacts with ADAM15 and FAS-L. Belonging to the sorting nexin family, SNX33 contains one BAR domain, one PX (phox homology) domain and one SH3 domain. The gene that encodes SNX33 consists of over 14,000 bases and maps to human chromosome 15q24.2. Housing approximately 106 million base pairs and encoding more than 700 genes, chromosome 15 makes up about 3% of the human genome. Angelman and Prader-Willi syndromes are associated with loss of function or deletion of genes in the 15q11-q13 region. In the case of Angelman syndrome, this loss is due to inactivity of the maternal 15q11-q13 encoded UBE3A gene in the brain by either chromosomal deletion or mutation. In cases of Prader-Willi syndrome, there is a partial or complete deletion of this region from the paternal copy of chromosome 15. Tay-Sachs disease is a lethal disorder associated with mutations of the HEXA gene, which is encoded by chromosome 15. Marfan syndrome is associated with chromosome 15 through the FBN1 gene.

REFERENCES

- Knoll, J.H., Nicholls, R.D., Magenis, R.E., Graham, J.M., Lalande, M. and Latt, S.A. 1989. Angelman and Prader-Willi syndromes share a common chromosome 15 deletion but differ in parental origin of the deletion. *Am. J. Med. Genet.* 32: 285-290.
- Hurowitz, G.I., Silver, J.M., Brin, M.F., Williams, D.T. and Johnson, W.G. 1993. Neuropsychiatric aspects of adult-onset Tay-Sachs disease: two case reports with several new findings. *J. Neuropsychiatry Clin. Neurosci.* 5: 30-36.
- Seet, L.F. and Hong, W. 2006. The Phox (PX) domain proteins and membrane traffic. *Biochim. Biophys. Acta* 1761: 878-896.
- Schöbel, S., Neumann, S., Hertweck, M., Dislich, B., Kuhn, P.H., Kremmer, E., Seed, B., Baumeister, R., Haass, C. and Lichtenthaler, S.F. 2008. A novel sorting nexin modulates endocytic trafficking and α -secretase cleavage of the amyloid precursor protein. *J. Biol. Chem.* 283: 14257-14268.
- Heiseke, A., Schöbel, S., Lichtenthaler, S.F., Vorberg, I., Groschup, M.H., Kretschmar, H., Schätzl, H.M. and Nunziante, M. 2008. The novel sorting nexin SNX33 interferes with cellular PrP formation by modulation of PrP shedding. *Traffic* 9: 1116-1129.
- Voss, M., Lettau, M. and Janssen, O. 2009. Identification of SH3 domain interaction partners of human FasL (CD178) by phage display screening. *BMC Immunol.* 10: 53.
- Zhang, J., Zhang, X., Guo, Y., Xu, L. and Pei, D. 2009. Sorting nexin 33 induces mammalian cell micronucleated phenotype and actin polymerization by interacting with Wiskott-Aldrich syndrome protein. *J. Biol. Chem.* 284: 21659-21669.
- Kleino, I., Ortiz, R.M., Yritys, M., Huovila, A.P. and Saksela, K. 2009. Alternative splicing of ADAM15 regulates its interactions with cellular SH3 proteins. *J. Cell. Biochem.* 108: 877-885.

CHROMOSOMAL LOCATION

Genetic locus: Snx33 (mouse) mapping to 9 B.

PRODUCT

SNX33 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SNX33 shRNA Plasmid (m): sc-153678-SH and SNX33 shRNA (m) Lentiviral Particles: sc-153678-V as alternate gene silencing products.

For independent verification of SNX33 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-153678A, sc-153678B and sc-153678C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SNX33 siRNA (m) is recommended for the inhibition of SNX33 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SNX33 gene expression knockdown using RT-PCR Primer: SNX33 (m)-PR: sc-153678-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.