

spectrin β IV siRNA (m): sc-153737

BACKGROUND

Spectrin, an Actin binding protein that is a major component of the cytoskeletal superstructure of the erythrocyte plasma membrane, is essential in determining the properties of the membrane including its shape and deformability. Spectrins function as membrane organizers and stabilizers, composed of non-homologous α and β chains, which aggregate side-to-side in an antiparallel fashion to form dimers, tetramers and higher polymers. The spectrin tetramers in erythrocytes act as barriers to lateral diffusion, but spectrin dimers seem to lack this function. Spectrin β IV is a non-erythrocytic member of the β -spectrin family. It is expressed in brain and pancreatic islets and localizes to the nuclear matrix, cytoplasmic vesicles and PML nuclear bodies. Spectrin β IV is a 2,564 amino acid protein with four isoforms due to alternative splicing events.

REFERENCES

1. Speicher, D.W. 1986. The present status of erythrocyte spectrin structure: the 106-residue repetitive structure is a basic feature of an entire class of proteins. *J. Cell. Biochem.* 30: 245-258.
2. Gardner, K., et al. 1987. Modulation of spectrin-actin assembly by erythrocyte adducin. *Nature* 328: 359-362.
3. Coleman, T.R., et al. 1989. Functional diversity among spectrin isoforms. *Cell Motil. Cytoskeleton* 12: 225-247.
4. Saxton, M.J. 1989. The spectrin network as a barrier to lateral diffusion in erythrocytes. A percolation analysis. *Biophys. J.* 55: 21-28.
5. Kennedy, S.P., et al. 1994. A partial structural repeat forms the heterodimer self-association site of all β -spectrins. *J. Biol. Chem.* 269: 11400-11408.
6. Nagase, T., et al. 2000. Prediction of the coding sequences of unidentified human genes. XVIII. The complete sequences of 100 new cDNA clones from brain which code for large proteins *in vitro*. *DNA Res.* 7: 273-281.
7. Tse, W.T., et al. 2001. A new spectrin β IV, has a major truncated isoform that associates with promyelocytic leukemia protein nuclear bodies and the nuclear matrix. *J. Biol. Chem.* 276: 23974-23985.
8. Boulanger, L., et al. 2002. Erythroid expression of the human α -spectrin gene promoter is mediated by GATA-1- and NF-E2-binding proteins. *J. Biol. Chem.* 277: 41563-41570.
9. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606214. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Sptbn4 (mouse) mapping to 7 A3.

PRODUCT

spectrin β IV siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see spectrin β IV shRNA Plasmid (m): sc-153737-SH and spectrin β IV shRNA (m) Lentiviral Particles: sc-153737-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

spectrin β IV siRNA (m) is recommended for the inhibition of spectrin β IV expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

spectrin β IV (F-5): sc-398506 is recommended as a control antibody for monitoring of spectrin β IV gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor spectrin β IV gene expression knockdown using RT-PCR Primer: spectrin β IV (m)-PR: sc-153737-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.