

Sptrx-3 siRNA (m): sc-153808

BACKGROUND

Sptrx-3 (spermatid-specific thioredoxin-3), also known as TRX6 (thioredoxin-6) or TXNDC8 (thioredoxin domain-containing protein 8), is a 127 amino acid protein that belongs to the thioredoxin family and contains one thioredoxin domain. Localizing to the cytoplasm and Golgi apparatus in testis, Sptrx-3 may be required for post-translational modifications of proteins required for acrosomal biogenesis and may also reduce disulfide bonds within the sperm. Sptrx-3 is only expressed during spermiogenesis, prominently in the Golgi apparatus of pachytene spermatocytes and round and elongated spermatids, with a transient localization in the developing acrosome of round spermatids. The Sptrx-3 gene contains 34,327 bases, encodes two alternatively spliced isoforms and maps to human chromosome 9q31.3. Human chromosome 9, which houses over 900 genes and comprises nearly 4% of the human genome, is associated with hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, and Familial dysautonomia.

REFERENCES

1. Miranda-Vizuete, A., Sadek, C.M., Jimenez, A., Krause, W.J., Sutovsky, P. and Oko, R. 2004. The mammalian testis-specific thioredoxin system. *Antioxid. Redox Signal.* 6: 25-40.
2. Jimenez, A., Zu, W., Rawe, V.Y., Pelto-Huikko, M., Flickinger, C.J., Sutovsky, P., Gustafsson, J.A., Oko, R. and Miranda-Vizuete, A. 2004. Spermatocyte/spermatid-specific thioredoxin-3, a novel Golgi apparatus-associated thioredoxin, is a specific marker of aberrant spermatogenesis. *J. Biol. Chem.* 279: 34971-34982.
3. Kondo, N., Nakamura, H., Masutani, H. and Yodoi, J. 2006. Redox regulation of human thioredoxin network. *Antioxid. Redox Signal.* 8: 1881-1890.
4. Cottin, V., Dupuis-Girod, S., Lesca, G. and Cordier, J.F. 2007. Pulmonary vascular manifestations of hereditary hemorrhagic telangiectasia (rendu-osler disease). *Respiration.* 74: 361-378.
5. Gold-von Simson, G., Goldberg, J.D., Rolnitzky, L.M., Mull, J., Leyne, M., Voustianiouk, A., Slangenaupt, S.A. and Axelrod, F.B. 2009. Kinetin in familial dysautonomia carriers: implications for a new therapeutic strategy targeting mRNA splicing. *Pediatr. Res.* 65: 341-346.
6. Buckman, C., George, T.C., Friend, S., Sutovsky, M., Miranda-Vizuete, A., Ozanon, C., Morrissey, P. and Sutovsky, P. 2009. High throughput, parallel imaging and biomarker quantification of human spermatozoa by ImageStream flow cytometry. *Syst. Biol. Reprod. Med.* 55: 244-251.
7. Axelrod, F.B., Hilz, M.J., Berlin, D., Yau, P.L., Javier, D., Sweat, V., Bruehl, H. and Convit, A. 2010. Neuroimaging supports central pathology in familial dysautonomia. *J. Neurol.* 257: 198-206.

CHROMOSOMAL LOCATION

Genetic locus: Txndc8 (mouse) mapping to 4 B3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Sptrx-3 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Sptrx-3 shRNA Plasmid (m): sc-153808-SH and Sptrx-3 shRNA (m) Lentiviral Particles: sc-153808-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Sptrx-3 siRNA (m) is recommended for the inhibition of Sptrx-3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Sptrx-3 gene expression knockdown using RT-PCR Primer: Sptrx-3 (m)-PR: sc-153808-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.