

Tektin 2 siRNA (m): sc-154182

BACKGROUND

Tektin 2 (TEKT2), also known as TEKTB1 (testicular tektin B1-like protein) or Tektin-t (testicular tektin), is a 430 amino acid protein that belongs to the Tektin family. While expressed at high levels in testis, trachea and fetal lung, Tektin 2 is expressed at lower levels in ovary, pituitary, adult lung, fetal brain and fetal kidney. Acting as the structural component of ciliary and flagellar microtubules, Tektin 2 forms filamentous polymers in the walls of those microtubules. Tektin 2 is tyrosine phosphorylated and plays a key role in the assembly or attachment of the inner dynein arm to microtubules in sperm flagella and tracheal cilia. The gene that encodes Tektin 2 consists of more than 4,000 bases and maps to human chromosome 1p34.3. Comprising nearly 8% of the human genome, chromosome 1 spans 260 million base pairs, contains over 3,000 genes and houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson's disease, Gaucher disease, schizophrenia and Usher syndrome.

REFERENCES

1. Eudy, J.D., Yao, S., Weston, M.D., Ma-Edmonds, M., Talmadge, C.B., Cheng, J.J., Kimberling, W.J. and Sumegi, J. 1998. Isolation of a gene encoding a novel member of the nuclear receptor superfamily from the critical region of Usher syndrome type IIa at 1q41. *Genomics* 50: 382-384.
2. Tayebi, N., Callahan, M., Madike, V., Stubblefield, B.K., Orvisky, E., Krasnewich, D., Fillano, J.J. and Sidransky, E. 2001. Gaucher disease and parkinsonism: a phenotypic and genotypic characterization. *Mol. Genet. Metab.* 73: 313-321.
3. Wolkowicz, M.J., Naaby-Hansen, S., Gamble, A.R., Reddi, P.P., Flickinger, C.J. and Herr, J.C. 2002. Tektin B1 demonstrates flagellar localization in human sperm. *Biol. Reprod.* 66: 241-250.
4. Iguchi, N., Tanaka, H., Nakamura, Y., Nozaki, M., Fujiwara, T. and Nishimune, Y. 2002. Cloning and characterization of the human Tektin-t gene. *Mol. Hum. Reprod.* 8: 525-530.
5. Plasilova, M., Russell, A.M., Wanner, A., Wolf, A., Dobbie, Z., Müller, H.J. and Heinemann, K. 2004. Exclusion of an extracolonic disease modifier locus on chromosome 1p33-36 in a large Swiss familial adenomatous polyposis kindred. *Eur. J. Hum. Genet.* 12: 365-371.
6. Tanaka, H., Iguchi, N., Toyama, Y., Kitamura, K., Takahashi, T., Kaseda, K., Maekawa, M. and Nishimune, Y. 2004. Mice deficient in the axonemal protein Tektin-t exhibit male infertility and immotile-cilium syndrome due to impaired inner arm dynein function. *Mol. Cell. Biol.* 24: 7958-7964.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 608953. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. Betarbet, R., Anderson, L.R., Gearing, M., Hodges, T.R., Fritz, J.J., Lah, J.J. and Levey, A.I. 2008. Fas-associated factor 1 and Parkinson's disease. *Neurobiol. Dis.* 31: 309-315.
9. Yokoi, T., Koide, R., Matsuoka, K., Nakagawa, A. and Azuma, N. 2009. Analysis of the vitreous membrane in a case of type 1 Stickler syndrome. *Graefes Arch. Clin. Exp. Ophthalmol.* 247: 715-718.

CHROMOSOMAL LOCATION

Genetic locus: Tekt2 (mouse) mapping to 4 D2.2.

PRODUCT

Tektin 2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Tektin 2 shRNA Plasmid (m): sc-154182-SH and Tektin 2 shRNA (m) Lentiviral Particles: sc-154182-V as alternate gene silencing products.

For independent verification of Tektin 2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154182A, sc-154182B and sc-154182C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Tektin 2 siRNA (m) is recommended for the inhibition of Tektin 2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Tektin 2 gene expression knockdown using RT-PCR Primer: Tektin 2 (m)-PR: sc-154182-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.