



TMED8 siRNA (m): sc-154336

BACKGROUND

TMED8 is a 325 amino acid protein that contains one GOLD domain. The gene that encodes TMED8 consists of approximately 35,361 bases and maps to human chromosome 14q24.3. Housing over 700 genes, chromosome 14 comprises nearly 3.5% of the human genome. Chromosome 14 encodes the Presenilin 1 (PSEN1) gene, which is one of the three key genes associated with the development of Alzheimer's disease. The SERPINA1 gene is also located on chromosome 14 and, when defective, leads to the genetic disorder α 1-antitrypsin deficiency, which is characterized by severe lung complications and liver dysfunction. An inversion of the long arm of chromosome 14 is thought to be involved in T-cell chronic lymphocytic leukemia (CLL), a cancer that affects T lymphocytes.

REFERENCES

1. Zech, L., Gahrton, G., Hammarström, L., Juliusson, G., Mellstedt, H., Robèrt, K.H. and Smith, C.I. 1984. Inversion of chromosome 14 marks human T-cell chronic lymphocytic leukaemia. *Nature* 308: 858-860.
2. Aisenberg, A.C., Krontiris, T.G., Mak, T.W. and Wilkes, B.M. 1985. Rearrangement of the gene for the β chain of the T-cell receptor in T-cell chronic lymphocytic leukemia and related disorders. *N. Engl. J. Med.* 313: 529-533.
3. Schellenberg, G.D., Bird, T.D., Wijsman, E.M., Orr, H.T., Anderson, L., Nemens, E., White, J.A., Bonycastle, L., Weber, J.L. and Alonso, M.E. 1992. Genetic linkage evidence for a familial Alzheimer's disease locus on chromosome 14. *Science* 258: 668-671.
4. Avramopoulos, D., Fallin, M.D. and Bassett, S.S. 2005. Linkage to chromosome 14q in Alzheimer's disease (AD) patients without psychotic symptoms. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 132B: 9-13.
5. Larner, A.J. and Doran, M. 2009. Genotype-phenotype relationships of presenilin-1 mutations in Alzheimer's disease: an update. *J. Alzheimers Dis.* 17: 259-265.
6. Topic, A., Alempijevic, T., Milutinovic, A.S. and Kovacevic, N. 2009. α 1-antitrypsin phenotypes in adult liver disease patients. *Ups. J. Med. Sci.* 114: 228-234.

CHROMOSOMAL LOCATION

Genetic locus: Tmed8 (mouse) mapping to 12 D2.

PRODUCT

TMED8 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TMED8 shRNA Plasmid (m): sc-154336-SH and TMED8 shRNA (m) Lentiviral Particles: sc-154336-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TMED8 siRNA (m) is recommended for the inhibition of TMED8 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TMED8 gene expression knockdown using RT-PCR Primer: TMED8 (m)-PR: sc-154336-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.