

TMPRSS5 siRNA (m): sc-154528

BACKGROUND

Extracellular proteases mediate the digestion of neighboring extracellular matrix components in initial tumor growth, allow desquamation of tumor cells into the surrounding environment, provide the basis for invasion of basement membranes in targeted metastatic organs and are required for release and activation of many growth and angiogenic factors. TMPRSS5 (transmembrane protease, serine 5), also known as spinesin, is a 457 amino acid single-pass type II membrane protein that is expressed specifically in brain and is thought to play a role in hearing. A member of the peptidase S1 family, TMPRSS5 contains one peptidase S1 domain and an SRCR domain, and is encoded by a gene that maps to human chromosome 11q23.2. Defects in the gene encoding TMPRSS5 are associated with deafness.

REFERENCES

1. Mohan, S., Thompson, G.R., Amaar, Y.G., Hathaway, G., Tschesche, H. and Baylink, D.J. 2002. ADAM-9 is an Insulin-like growth factor binding protein-5 protease produced and secreted by human osteoblasts. *Biochemistry* 41: 15394-15403.
2. Yamaguchi, N., Okui, A., Yamada, T., Nakazato, H. and Mitsui, S. 2002. Spinesin/TMPRSS5, a novel transmembrane serine protease, cloned from human spinal cord. *J. Biol. Chem.* 277: 6806-6812.
3. Guipponi, M., Toh, M.Y., Tan, J., Park, D., Hanson, K., Ballana, E., Kwong, D., Cannon, P.Z., Wu, Q., Gout, A., Delorenzi, M., Speed, T.P., Smith, R.J., Dahl, H.H., Petersen, M., Teasdale, R.D., Estivill, X., Park, W.J., et al. 2008. An integrated genetic and functional analysis of the role of type II transmembrane serine proteases (TMPRSSs) in hearing loss. *Hum. Mutat.* 29: 130-141.
4. Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 606751. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Bugge, T.H., Antalis, T.M. and Wu, Q. 2009. Type II transmembrane serine proteases. *J. Biol. Chem.* 284: 23177-23181.
6. Ha, Y. 2009. Structure and mechanism of intramembrane protease. *Semin. Cell Dev. Biol.* 20: 240-250.

CHROMOSOMAL LOCATION

Genetic locus: *Tmprss5* (mouse) mapping to 9 A5.3.

PRODUCT

TMPRSS5 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TMPRSS5 shRNA Plasmid (m): sc-154528-SH and TMPRSS5 shRNA (m) Lentiviral Particles: sc-154528-V as alternate gene silencing products.

For independent verification of TMPRSS5 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154528A, sc-154528B and sc-154528C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TMPRSS5 siRNA (m) is recommended for the inhibition of TMPRSS5 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TMPRSS5 gene expression knockdown using RT-PCR Primer: TMPRSS5 (m)-PR: sc-154528-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.