

TXNDC9 siRNA (m): sc-154824

BACKGROUND

Thioredoxins comprise a family of small proteins that, by catalyzing the oxidation of disulfide bonds, participate in redox reactions throughout the cell. Proteins that contain thioredoxin domains do not necessarily convey the oxidative properties of thioredoxins, but generally function as disulfide isomerases that enzymatically rearrange disulfide bonds found in various proteins. TXNDC9 (thioredoxin domain-containing protein 9), also known as APACD (ATP-binding protein associated with cell differentiation), is a 226 amino acid protein that contains one thioredoxin domain and may be involved in cell differentiation events. The gene encoding TXNDC9 maps to human chromosome 2, which houses over 1,400 genes and comprises nearly 8% of the human genome. Harlequin ichthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene, while the lipid metabolic disorder sitosterolemia is associated with defects in the ABCG5 and ABCG8 genes. Additionally, an extremely rare recessive genetic disorder, Alström syndrome, is caused by mutations in the ALMS1 gene, which maps to chromosome 2.

REFERENCES

- Holmgren, A. 1985. Thioredoxin. *Annu. Rev. Biochem.* 54: 237-271.
- Holmgren, A. 1989. Thioredoxin and glutaredoxin systems. *J. Biol. Chem.* 264: 13963-13966.
- Eklund, H., et al. 1991. Structural and functional relations among thioredoxins of different species. *Proteins* 11: 13-28.
- Ijdo, J.W., et al. 1991. Origin of human chromosome 2: an ancestral telomere-telomere fusion. *Proc. Natl. Acad. Sci. USA* 88: 9051-9055.
- Freedman, R.B., et al. 1994. Protein disulphide isomerase: building bridges in protein folding. *Trends Biochem. Sci.* 19: 331-336.
- Ogawa, S., et al. 2004. An evolutionarily conserved gene required for proper microtubule architecture in *Caenorhabditis elegans*. *Genes Cells* 9: 83-93.
- Thomas, A.C., et al. 2006. ABCA12 is the major harlequin ichthyosis gene. *J. Invest. Dermatol.* 126: 2408-2413.
- Marshall, J.D., et al. 2007. Alström syndrome. *Eur. J. Hum. Genet.* 15: 1193-1202.

CHROMOSOMAL LOCATION

Genetic locus: Txnnc9 (mouse) mapping to 1 B.

PRODUCT

TXNDC9 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TXNDC9 shRNA Plasmid (m): sc-154824-SH and TXNDC9 shRNA (m) Lentiviral Particles: sc-154824-V as alternate gene silencing products.

For independent verification of TXNDC9 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154824A, sc-154824B and sc-154824C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TXNDC9 siRNA (m) is recommended for the inhibition of TXNDC9 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

TXNDC9 (C-8): sc-514770 is recommended as a control antibody for monitoring of TXNDC9 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TXNDC9 gene expression knockdown using RT-PCR Primer: TXNDC9 (m)-PR: sc-154824-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.