



U11/U12 snRNP 35K siRNA (m): sc-154832

BACKGROUND

Small nuclear ribonucleoproteins, also known as snRNPs, combine with other proteins to form spliceosomes, a complex that catalyzes pre-mRNA splicing. There are two types of spliceosomes: U2 and U12. The U2-type spliceosome is found in all eukaryotes and excises U2-type introns, which account for the majority of pre-mRNA introns. The U12-type spliceosome removes U12-type introns, which comprise less than 1% of all human introns. The U12-type spliceosome is comprised of the U11 and U12 snRNPs as well as the U4/U6.U5 tri-snRNP. U11 and U12 bind as a U11/U12 di-snRNP complex, which recognizes the 5' splice site of the pre-mRNA during the first steps of U12-type spliceosome formation. U11/U12 snRNPs contain several proteins, including seven that are unique to the U11/U12snRNP: 65K, 59K, 48K, 35K, 31K, 25K and 20K. U11/U12 snRNP 35K is a 246 amino acid protein localized to the nucleus, and it contains a motif known to mediate RNA binding. Two named isoforms of this protein exist as a result of alternative splicing events.

REFERENCES

- Will, C.L., Schneider, C., MacMillan, A.M., Katopodis, N.F., Neubauer, G., Wilm, M., Lührmann, R. and Query, C.C. 2001. A novel U2 and U11/U12 snRNP protein that associates with the pre-mRNA branch site. *EMBO J.* 20: 4536-4546.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 610750. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Will, C.L., Schneider, C., Hossbach, M., Urlaub, H., Rahut, R., Elbashir, S., Tuschl, T. and Lührmann, R. 2004. The human 18S U11/U12 snRNP contains a set of novel proteins not found in the U2-dependent spliceosome. *RNA* 10: 929-941.
- Benecke, H., Lührmann, R. and Will, C.L. 2005. The U11/U12 snRNP 65K protein acts as a molecular bridge, binding the U12 snRNA and U11-59K protein. *EMBO J.* 24: 3057-3069.
- Golas, M.M., Sander, B., Will, C.L., Lührmann, R. and Stark, H. 2005. Major conformational change in the complex SF3b upon integration into the spliceosomal U11/U12 di-snRNP as revealed by electron cryomicroscopy. *Mol. Cell* 17: 869-883.
- Schellenberg, M.J., Edwards, R.A., Ritchie, D.B., Kent, O.A., Golas, M.M., Stark, H., Lührmann, R., Glover, J.N. and MacMillan, A.M. 2006. Crystal structure of a core spliceosomal protein interface. *Proc. Natl. Acad. Sci. USA* 103: 1266-1271.
- Wang, H., Gao, M.X., Li, L., Wang, B., Hori, N. and Sato, K. 2007. Isolation, expression, and characterization of the human ZCRB1 gene mapped to 12q12. *Genomics* 89: 59-69.
- Turunen, J.J., Will, C.L., Grote, M., Lührmann, R. and Frilander, M.J. 2008. The U11-48K protein contacts the 5' splice site of U12-type introns and the U11-59K protein. *Mol. Cell. Biol.* 28: 3548-3560.

CHROMOSOMAL LOCATION

Genetic locus: *Snrnp35* (mouse) mapping to 5 F.

PRODUCT

U11/U12 snRNP 35K siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see U11/U12 snRNP 35K shRNA Plasmid (m): sc-154832-SH and U11/U12 snRNP 35K shRNA (m) Lentiviral Particles: sc-154832-V as alternate gene silencing products.

For independent verification of U11/U12 snRNP 35K (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154832A, sc-154832B and sc-154832C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

U11/U12 snRNP 35K siRNA (m) is recommended for the inhibition of U11/U12 snRNP 35K expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor U11/U12 snRNP 35K gene expression knockdown using RT-PCR Primer: U11/U12 snRNP 35K (m)-PR: sc-154832-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.