



UCRC siRNA (m): sc-154888

BACKGROUND

UCRC (cytochrome b-c1 complex subunit 9), also known as ubiquinol-cytochrome c reductase complex 7.2 kDa protein and complex III subunit X (UQCR10), is a 63 amino acid mitochondrial inner membrane protein that is a component of the ubiquinol-cytochrome c reductase complex, a part of the mitochondrial respiratory chain. This complex consists of eleven subunits and interacts with cytochrome c1. Ubiquinol cytochrome c reductase (complex III) deficiency results when any of the genes encoding the subunits is missing or defective, and causes cardiomyopathy, Leber's myopathy and fatal infant conditions. UCRC is highly expressed in skeletal muscle and heart, where its levels have been shown to be influenced by ovarian hormones. Interestingly, UCRC expression in the heart increases in response to cardiac hypertrophy and hypertension.

REFERENCES

1. Reed, J. and Hess, B. 1977. Structure of a cytochrome b-c 1 complex from *Saccharomyces cerevisiae* YF. Hoppe-Seyler's Z. Physiol. Chem. 358: 1119-1124.
2. Schägger, H., Brandt, U., Gencic, S. and von Jagow, G. 1995. Ubiquinol-cytochrome-c reductase from human and bovine mitochondria. Meth. Enzymol. 260: 82-96.
3. Islam, M.M., Suzuki, H., Yoneda, M. and Tanaka, M. 1997. Primary structure of the smallest (6.4-kDa) subunit of human and bovine ubiquinol-cytochrome c reductase deduced from cDNA sequences. Biochem. Mol. Biol. Int. 41: 1109-1116.
4. Valnot, I., Kassiss, J., Chretien, D., de Lonlay, P., Parfait, B., Munnich, A., Kachaner, J., Rustin, P. and Rötig, A. 1999. A mitochondrial cytochrome b mutation but no mutations of nuclearly encoded subunits in ubiquinol cytochrome c reductase (complex III) deficiency. Hum. Genet. 104: 460-466.
5. Zhang, Q.H., Ye, M., Wu, X.Y., Ren, S.X., Zhao, M., Zhao, C.J., Fu, G., Shen, Y., Fan, H.Y., Lu, G., Zhong, M., Xu, X.R., Han, Z.G., Zhang, J.W., Tao, J., Huang, Q.H., Zhou, J., Hu, G.X., Gu, J., Chen, S.J. and Chen, Z. 2000. Cloning and functional analysis of cDNAs with open reading frames for 300 previously undefined genes expressed in CD34⁺ hematopoietic stem/progenitor cells. Genome Res. 10: 1546-1560.
6. Akashi, H., Han, H.J., Iizaka, M., Nakajima, Y., Furukawa, Y., Sugano, S., Imai, K. and Nakamura, Y. 2000. Isolation and characterization of a human cDNA encoding a protein homologous to the 7.2-kDa protein (subunit X) of bovine ubiquinol-cytochrome C reductase. J. Hum. Genet. 45: 43-46.
7. Huynh, H., Servant, N. and Chalifour, L.E. 2007. Ubiquinol-cytochrome-c reductase 7.2 kDa protein of mitochondrial complex III is steroid-responsive and increases in cardiac hypertrophy and hypertension. Can. J. Physiol. Pharmacol. 85: 986-996.
8. Yokota, T., Kinugawa, S., Hirabayashi, K., Matsushima, S., Inoue, N., Ohta, Y., Hamaguchi, S., Sobirin, M.A., Ono, T., Suga, T., Kuroda, S., Tanaka, S., Terasaki, F., Okita, K. and Tsutsui, H. 2009. Oxidative stress in skeletal muscle impairs mitochondrial respiration and limits exercise capacity in type 2 diabetic mice. Am. J. Physiol. Heart Circ. Physiol. 297: H1069-H1077.

CHROMOSOMAL LOCATION

Genetic locus: Uqcr10 (mouse) mapping to 11 A1.

PRODUCT

UCRC siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see UCRC shRNA Plasmid (m): sc-154888-SH and UCRC shRNA (m) Lentiviral Particles: sc-154888-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

UCRC siRNA (m) is recommended for the inhibition of UCRC expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.