

WBSCR18 siRNA (m): sc-155247

BACKGROUND

Williams-Beuren syndrome (WBS) is a developmental disorder caused by the hemizygous microdeletion on chromosome 7q11.23. WBS is an autosomal dominant genetic condition that is characterized by physical, cognitive and behavioral traits. The physical traits associated with WBS include facial dysmorphism, vascular stenoses, growth deficiencies, dental anomalies and neurologic and musculoskeletal abnormalities. Mild retardation, a weakness in visual-spatial skills, anxiety and a short attention span are typical cognitive and behavioral traits of WBS patients. The WBSCR18 gene is located within the WBS deletion and may contribute to the developmental symptoms found in WBS because of a loss of the encoded transcription factor. WBSCR18 (Williams-Beuren syndrome chromosomal region 18 protein) is a 226 amino acid protein containing one J domain. WBSCR18 is expressed in brain, heart, kidney, liver, lung, spleen, stomach and testis.

REFERENCES

1. Morris, C.A., et al. 1988. Natural history of Williams syndrome: physical characteristics. *J. Pediatr.* 113: 318-326.
2. Pober, B.R. and Dykens, E.M. 1996. Williams syndrome: an overview of medical, cognitive, and behavioral features. *Child Adolesc. Psychiatr. Clin. North Am.* 5: 929-943.
3. Lashkari, A., et al. 1999. Williams-Beuren syndrome: an update and review for the primary physician. *Clin. Pediatr.* 38: 189-208.
4. Bellugi, U., et al. 1999. Bridging cognition, the brain and molecular genetics: evidence from Williams syndrome. *Trends Neurosci.* 22: 197-207.
5. Merla, G., et al. 2002. Identification of additional transcripts in the Williams-Beuren syndrome critical region. *Hum. Genet.* 110: 429-438.
6. Hillier, L.W., et al. 2003. The DNA sequence of human chromosome 7. *Nature* 424: 157-164.
7. Ota, T., et al. 2004. Complete sequencing and characterization of 21,243 full-length human cDNAs. *Nat. Genet.* 36: 40-45.
8. Gerhard, D.S., et al. 2004. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). *Genome Res.* 14: 2121-2127.

CHROMOSOMAL LOCATION

Genetic locus: Dnajc30 (mouse) mapping to 5 G2.

PRODUCT

WBSCR18 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see WBSCR18 shRNA Plasmid (m): sc-155247-SH and WBSCR18 shRNA (m) Lentiviral Particles: sc-155247-V as alternate gene silencing products.

For independent verification of WBSCR18 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155247A, sc-155247B and sc-155247C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

WBSCR18 siRNA (m) is recommended for the inhibition of WBSCR18 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor WBSCR18 gene expression knockdown using RT-PCR Primer: WBSCR18 (m)-PR: sc-155247-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.