

WFDC12 siRNA (m): sc-155336

BACKGROUND

Peptidases are enzymes that are responsible for hydrolyzing peptide bonds of polypeptide chains during protein catabolism. Protease inhibitors are important peptidase regulators which halt enzymatic function. The WAP (whey acidic protein) domain, also referred to as the WAP-type four-disulfide core domain, is a signature protein motif that contains eight cysteine residues which form disulfide bonds and may exhibit protease inhibitor activity. WAP domain-containing proteins are thought to function in the immune defense by cleaving microbial proteolytic enzymes in order to prevent tissue penetration and infection. WFDC12 (WAP four-disulfide core domain protein 12), also known as WAP2, is a 111 amino acid secreted protein that contains one WAP domain and is highly expressed in lung, esophagus, prostate and skin. A cluster of WAP genes, including WFDC12, exist on chromosome 20, suggesting they evolved by repeated duplications.

REFERENCES

1. Brem, G., Brenig, B., Salmons, B., Wolf, E., Müller, M., Erfle, V., Günzburg, W.H. and Dahme, E. 1991. Unexpected transgene expression of a mammary-specific growth hormone gene construct in Bergmann glial cells of the mouse. *Tierarztl. Prax.* 19: 345-350.
2. Aigner, B., Pambalk, K., Reichart, U., Besenfelder, U., Bosze, Z., Renner, M., Günzburg, W.H., Wolf, E., Müller, M. and Brem, G. 1999. Species-specific alternative splicing of transgenic RNA in the mammary glands of pigs, rabbits, and mice. *Biochem. Biophys. Res. Commun.* 257: 843-850.
3. Lundwall, A. and Clauss, A. 2002. Identification of a novel protease inhibitor gene that is highly expressed in the prostate. *Biochem. Biophys. Res. Commun.* 290: 452-456.
4. Clauss, A., Lilja, H. and Lundwall, A. 2002. A locus on human chromosome 20 contains several genes expressing protease inhibitor domains with homology to whey acidic protein. *Biochem. J.* 368: 233-242.
5. Clark, H.F., Gurney, A.L., Abaya, E., Baker, K., Baldwin, D., Brush, J., Chen, J., Chow, B., Chui, C., Crowley, C., Currell, B., Deuel, B., Dowd, P., Eaton, D., Foster, J., Grimaldi, C., Gu, Q., Hass, P.E., Heldens, S., Huang, A., et al. 2003. The secreted protein discovery initiative (SPDI), a large-scale effort to identify novel human secreted and transmembrane proteins: a bioinformatics assessment. *Genome Res.* 13: 2265-2270.
6. Clauss, A., Lilja, H. and Lundwall, A. 2005. The evolution of a genetic locus encoding small serine proteinase inhibitors. *Biochem. Biophys. Res. Commun.* 333: 383-389.
7. Lundwall, A. 2007. A locus on chromosome 20 encompassing genes that are highly expressed in the epididymis. *Asian J. Androl.* 9: 540-544.
8. Hurle, B., Swanson, W. and Green, E.D. 2007. Comparative sequence analyses reveal rapid and divergent evolutionary changes of the WFDC locus in the primate lineage. *Genome Res.* 17: 276-286.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Wfdc12 (mouse) mapping to 2 H3.

PRODUCT

WFDC12 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see WFDC12 shRNA Plasmid (m): sc-155336-SH and WFDC12 shRNA (m) Lentiviral Particles: sc-155336-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

WFDC12 siRNA (m) is recommended for the inhibition of WFDC12 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor WFDC12 gene expression knockdown using RT-PCR Primer: WFDC12 (m)-PR: sc-155336-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.