



WSB2 siRNA (m): sc-155362

BACKGROUND

WD-repeats are motifs that are found in a variety of proteins and are characterized by a conserved core of 40-60 amino acids that commonly form a tertiary propeller structure. While proteins that contain WD-repeats participate in a wide range of cellular functions, they are generally involved in regulatory mechanisms concerning chromatin assembly, cell cycle control, signal transduction, RNA processing, apoptosis and vesicular trafficking. WSB2 (WD repeat and SOCS box-containing 2), also known as CS box-containing WD protein or SBA2, is a 404 amino acid protein containing five WD repeats and one SOCS box domain, which may function as a bridge between certain substrate-binding domains and E3 ubiquitin protein ligases. The gene encoding WSB2 maps to human chromosome 12, which encodes over 1,100 genes, comprises approximately 4.5% of the human genome and is associated with a variety of diseases and afflictions including hypochondrogenesis, achondrogenesis, Kniest dysplasia and Noonan syndrome.

REFERENCES

1. Garcia-Higuera, I., et al. 1998. Folding a WD repeat propeller. Role of highly conserved aspartic acid residues in the G protein β subunit and Sec13. *J. Biol. Chem.* 273: 9041-9049.
2. Delgado Carrasco, J., et al. 2001. Achondrogenesis type II-hypochondrogenesis: radiological features. *Case report. An. Esp. Pediatr.* 55: 553-557.
3. Kile, B.T. and Alexander, W.S. 2001. The suppressors of cytokine signalling (SOCS). *Cell. Mol. Life Sci.* 58: 1627-1635.
4. Li, D. and Roberts, R. 2001. WD-repeat proteins: structure characteristics, biological function, and their involvement in human diseases. *Cell. Mol. Life Sci.* 58: 2085-2097.
5. Kile, B.T., et al. 2002. The SOCS box: a tale of destruction and degradation. *Trends Biochem. Sci.* 27: 235-241.
6. Yokoyama, T., et al. 2003. A case of Kniest dysplasia with retinal detachment and the mutation analysis. *Am. J. Ophthalmol.* 136: 1186-1188.

CHROMOSOMAL LOCATION

Genetic locus: Wsb2 (mouse) mapping to 5 F.

PRODUCT

WSB2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see WSB2 shRNA Plasmid (m): sc-155362-SH and WSB2 shRNA (m) Lentiviral Particles: sc-155362-V as alternate gene silencing products.

For independent verification of WSB2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155362A, sc-155362B and sc-155362C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

WSB2 siRNA (m) is recommended for the inhibition of WSB2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor WSB2 gene expression knockdown using RT-PCR Primer: WSB2 (m)-PR: sc-155362-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.