

ZFP1 siRNA (m): sc-155519

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZFP1 (zinc finger protein 1), also known as ZNF475, is a 407 amino acid protein that contains one KRAB domain and eight C₂H₂-type zinc fingers. Localizes to the nucleus, ZFP1 exists as multiple alternatively spliced isoforms and is thought to play a role in transcriptional regulation events. The gene encoding ZFP1 maps to human chromosome 16q23.1, which encodes over 900 genes and comprises nearly 3% of the human genome.

REFERENCES

1. Chowdhury, K., Rohdewohld, H. and Gruss, P. 1988. Specific and ubiquitous expression of different Zn finger protein genes in the mouse. *Nucleic Acids Res.* 16: 9995-10011.
2. Chowdhury, K., Dietrich, S., Balling, R., Guenet, J.L. and Gruss, P. 1989. Structure, expression and chromosomal localization of Zfp-1, a murine zinc finger protein gene. *Nucleic Acids Res.* 17: 10427-10438.
3. South, T.L., Kim, B., Hare, D.R. and Summers, M.F. 1990. Zinc fingers and molecular recognition. Structure and nucleic acid binding studies of an HIV zinc finger-like domain. *Biochem. Pharmacol.* 40: 123-129.
4. Gilbert, F. 1999. Disease genes and chromosomes: disease maps of the human genome. *Chromosome 16. Genet. Test.* 3: 243-254.
5. Sun, Y., Gou, D.M., Liu, H., Peng, X. and Li, W.X. 2003. The KRAB domain of zinc finger gene ZNF268: a potential transcriptional repressor. *IUBMB Life* 55: 127-131.
6. O'Geen, H., Squazzo, S.L., Iyengar, S., Blahnik, K., Rinn, J.L., Chang, H.Y., Green, R. and Farnham, P.J. 2007. Genome-wide analysis of KAP1 binding suggests autoregulation of KRAB-ZNFs. *PLoS Genet.* 3: e89.

CHROMOSOMAL LOCATION

Genetic locus: Zfp1 (mouse) mapping to 8 E1.

PRODUCT

ZFP1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZFP1 shRNA Plasmid (m): sc-155519-SH and ZFP1 shRNA (m) Lentiviral Particles: sc-155519-V as alternate gene silencing products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZFP1 siRNA (m) is recommended for the inhibition of ZFP1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZFP1 gene expression knockdown using RT-PCR Primer: ZFP1 (m)-PR: sc-155519-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.