

ZFYVE21 siRNA (m): sc-155605

BACKGROUND

ZFYVE21 (zinc finger FYVE domain-containing protein 21) is a 234 amino acid protein that localizes to cell junctions and has one FYVE-type zinc finger. ZFYVE21 exists as two isoforms generated by alternative splicing events. The gene encoding ZFYVE1 maps to human chromosome 14q24.2, which houses over 700 genes and comprises nearly 3.5% of the human genome. Chromosome 14 encodes the presenilin 1 (PSEN1) gene, which is one of the three key genes associated with the development of Alzheimer's disease (AD). The SERPINA1 gene is also located on chromosome 14 and, when defective, leads to the genetic disorder α 1-antitrypsin deficiency, which is characterized by severe lung complications and liver dysfunction.

REFERENCES

1. Ridley, S.H., Ktistakis, N., Davidson, K., Anderson, K.E., Manifava, M., Ellson, C.D., Lipp, P., Bootman, M., Coadwell, J., Nazarian, A., Erdjument-Bromage, H., Tempst, P., Cooper, M.A., Thuring, J.W., et al. 2001. FENS-1 and DFCEP1 are FYVE domain-containing proteins with distinct functions in the endosomal and Golgi compartments. *J. Cell Sci.* 114: 3991-4000.
2. Avramopoulos, D., Fallin, M.D. and Bassett, S.S. 2005. Linkage to chromosome 14q in Alzheimer's disease (AD) patients without psychotic symptoms. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 132B: 9-13.
3. Larner, A.J. and Doran, M. 2009. Genotype-phenotype relationships of presenilin-1 mutations in Alzheimer's disease: an update. *J. Alzheimers Dis.* 17: 259-265.
4. Topic, A., Alempijevic, T., Milutinovic, A.S. and Kovacevic, N. 2009. α 1-antitrypsin phenotypes in adult liver disease patients. *Ups. J. Med. Sci.* 114: 228-234.
5. Nagano, M., Hoshino, D., Sakamoto, T., Kawasaki, N., Koshikawa, N. and Seiki, M. 2010. ZF21 protein regulates cell adhesion and motility. *J. Biol. Chem.* 285: 21013-21022.

CHROMOSOMAL LOCATION

Genetic locus: Zfyve21 (mouse) mapping to 12 F1.

PRODUCT

ZFYVE21 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZFYVE21 shRNA Plasmid (m): sc-155605-SH and ZFYVE21 shRNA (m) Lentiviral Particles: sc-155605-V as alternate gene silencing products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZFYVE21 siRNA (m) is recommended for the inhibition of ZFYVE21 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZFYVE21 gene expression knockdown using RT-PCR Primer: ZFYVE21 (m)-PR: sc-155605-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.