



## Zic4 siRNA (m): sc-155611

### BACKGROUND

Zic4 (zinc finger protein of the cerebellum 4) is a C<sub>2</sub>H<sub>2</sub> zinc finger transcription factor that influences cerebellar development. Zic4 localizes to the nuclei of cerebellar granule cells. Zic4 mRNA expression peaks on postnatal day 5 in the developing cerebellum of mouse. Zic family members are important during development, and have been associated with X-linked visceral heterotaxy and holoprosencephaly type 5. Zic4 is closely linked to Zic1, a related family member on chromosome 3q24.

### REFERENCES

1. Aruga, J., et al. 1996. Identification and characterization of Zic4, a new member of the mouse Zic gene family. *Gene* 172: 291-294.
2. Nagai, T., et al. 1997. The expression of the mouse Zic1, Zic2, and Zic3 gene suggests an essential role for Zic genes in body pattern formation. *Dev. Biol.* 182: 299-313.
3. Ogura, H., et al. 2001. Behavioral abnormalities of Zic1 and Zic2 mutant mice: implications as models for human neurological disorders. *Behav. Genet.* 31: 317-324.
4. Salero, E., et al. 2001. Transcription factors Zic1 and Zic2 bind and transactivate the apolipoprotein E gene promoter. *J. Biol. Chem.* 276: 1881-1888.
5. Ebert, P.J., et al. 2003. Zic1 represses Math1 expression via interactions with the Math1 enhancer and modulation of Math1 autoregulation. *Development* 130: 1949-1959.
6. Grinberg, I., et al. 2004. Heterozygous deletion of the linked genes ZIC1 and ZIC4 is involved in Dandy-Walker malformation. *Nat. Genet.* 36: 1053-1055.
7. LocusLink Report (LocusID: 84107). <http://www.ncbi.nlm.nih.gov/LocusLink/>
8. <http://harvester.embl.de/harvester/Q8N9/Q8N9L1.htm>

### CHROMOSOMAL LOCATION

Genetic locus: Zic4 (mouse) mapping to 9 E3.3.

### PRODUCT

Zic4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Zic4 shRNA Plasmid (m): sc-155611-SH and Zic4 shRNA (m) Lentiviral Particles: sc-155611-V as alternate gene silencing products.

For independent verification of Zic4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155611A, sc-155611B and sc-155611C.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

Zic4 siRNA (m) is recommended for the inhibition of Zic4 expression in mouse cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Zic4 gene expression knockdown using RT-PCR Primer: Zic4 (m)-PR: sc-155611-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.