

ZNF114 siRNA (m): sc-155636

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF114 (zinc finger protein 114) is a 417 amino acid nuclear protein that is implicated in transcriptional regulation and exists as two alternatively spliced isoforms. A member of the Krüppel C₂H₂-type zinc-finger protein family, ZNF114 contains four C₂H₂-type zinc fingers and one KRAB domain, and is encoded by a gene that maps to human chromosome 19q13.33. Human chromosome 19 consists of over 63 million bases, houses approximately 1,400 genes and is recognized for having the greatest gene density of the human chromosomes.

REFERENCES

1. Bray, P., Lichter, P., Thiesen, H.J., Ward, D.C. and Dawid, I.B. 1991. Characterization and mapping of human genes encoding zinc finger proteins. *Proc. Natl. Acad. Sci. USA* 88: 9563-9567.
2. Lichter, P., Bray, P., Ried, T., Dawid, I.B. and Ward, D.C. 1992. Clustering of C₂H₂ zinc finger motif sequences within telomeric and fragile site regions of human chromosomes. *Genomics* 13: 999-1007.
3. Bellefroid, E.J., Marine, J.C., Ried, T., Lecocq, P.J., Rivière, M., Amemiya, C., Poncelet, D.A., Coulie, P.G., de Jong, P. and Szpirer, C. 1993. Clustered organization of homologous KRAB zinc-finger genes with enhanced expression in human T lymphoid cells. *EMBO J.* 12: 1363-1374.
4. Dehal, P., Predki, P., Olsen, A.S., Kobayashi, A., Foltz, P., Lucas, S., Land, M., Terry, A., Ecale Zhou, C.L., Rash, S., Zhang, Q., Gordon, L., Kim, J., Elkin, C., Pollard, M.J., Richardson, P., Rokhsar, D., Uberbacher, E., Hawkins, T., et al. 2001. Human chromosome 19 and related regions in mouse: conservative and lineage-specific evolution. *Science* 293: 104-111.
5. Looman, C., Hellman, L. and Abrink, M. 2004. A novel Krüppel-associated box identified in a panel of mammalian zinc finger proteins. *Mamm. Genome* 15: 35-40.
6. Huntley, S., Baggott, D.M., Hamilton, A.T., Tran-Gyamfi, M., Yang, S., Kim, J., Gordon, L., Branscomb, E. and Stubbs, L. 2006. A comprehensive catalog of human KRAB-associated zinc finger genes: insights into the evolutionary history of a large family of transcriptional repressors. *Genome Res.* 16: 669-677.
7. Filion, G.J., Zhenilo, S., Salozhin, S., Yamada, D., Prokhortchouk, E. and Defossez, P.A. 2006. A family of human zinc finger proteins that bind methylated DNA and repress transcription. *Mol. Cell. Biol.* 26: 169-181.
8. Tian, C.Y., Zhang, L.Q. and He, F.C. 2006. Progress in the study of KRAB zinc finger protein. *Yi Chuan* 28: 1451-1456.

CHROMOSOMAL LOCATION

Genetic locus: Zfp114 (mouse) mapping to 7 A3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

ZNF114 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF114 shRNA Plasmid (m): sc-155636-SH and ZNF114 shRNA (m) Lentiviral Particles: sc-155636-V as alternate gene silencing products.

For independent verification of ZNF114 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155636A, sc-155636B and sc-155636C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZNF114 siRNA (m) is recommended for the inhibition of ZNF114 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF114 gene expression knockdown using RT-PCR Primer: ZNF114 (m)-PR: sc-155636-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.