

ZNF239 siRNA (m): sc-155664

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF239, also known as zinc finger protein MOK-2 or HOK-2, is a 458 amino acid protein belonging to the Krüppel C₂H₂-type zinc-finger protein family. Localized to the nucleus, ZNF239 contains nine C₂H₂-type zinc finger domains. Due to the presence of these domains, ZNF239 may be involved in transcriptional regulation. ZNF239 is present at high levels in brain, breast and testis, and has no expression in liver or placenta.

REFERENCES

1. Bellefroid, E.J., Poncelet, D.A., Lecocq, P.J., Revelant, O. and Martial, J.A. 1991. The evolutionarily conserved Krüppel-associated box domain defines a subfamily of eukaryotic multifingered proteins. *Proc. Natl. Acad. Sci. USA* 88: 3608-3612.
2. Margolin, J.F., Friedman, J.R., Meyer, W.K., Vissing, H., Thiesen, H.J. and Rauscher, F.J. 1994. Krüppel-associated boxes are potent transcriptional repression domains. *Proc. Natl. Acad. Sci. USA* 91: 4509-4513.
3. Ernault-Lange, M., Arranz, V., Le Coniat, M., Berger, R. and Kress, M. 1995. Human and mouse Krüppel-like (MOK2) orthologue genes encode two different zinc finger proteins. *J. Mol. Evol.* 41: 784-794.
4. Arranz, V., Kress, M. and Ernault-Lange, M. 1996. Localization of zinc finger Mok2 gene to mouse chromosome 6, a new region of homology with human chromosome 19. *Mamm. Genome* 7: 77-78.
5. Arranz, V., Harper, F., Florentin, Y., Puvion, E., Kress, M. and Ernault-Lange, M. 1997. Human and mouse MOK2 proteins are associated with nuclear ribonucleoprotein components and bind specifically to RNA and DNA through their zinc finger domains. *Mol. Cell. Biol.* 17: 2116-2126.
6. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 601069. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Shannon, M., Hamilton, A.T., Gordon, L., Branscomb, E. and Stubbs, L. 2003. Differential expansion of zinc-finger transcription factor loci in homologous human and mouse gene clusters. *Genome Res.* 13: 1097-1110.
8. Dreuillet, C., Harper, M., Tillit, J., Kress, M. and Ernault-Lange, M. 2008. Mislocalization of human transcription factor MOK2 in the presence of pathogenic mutations of Lamin A/C. *Biol. Cell* 100: 51-61.

CHROMOSOMAL LOCATION

Genetic locus: Zfp239 (mouse) mapping to 6 F1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

ZNF239 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF239 shRNA Plasmid (m): sc-155664-SH and ZNF239 shRNA (m) Lentiviral Particles: sc-155664-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZNF239 siRNA (m) is recommended for the inhibition of ZNF239 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF239 gene expression knockdown using RT-PCR Primer: ZNF239 (m)-PR: sc-155664-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.