

# zygin 2 siRNA (m): sc-155855

## BACKGROUND

Zygin 2, also known as FEZ2 (fasciculation and elongation protein  $\zeta$  2), is a 353 amino acid cytoplasmic protein that belongs to the zygin family. Zygin 2 exists as a difulvide-linked homodimer but may form heterodimers with zygin 1. Expressed in non-neural tissues including heart, lung, spleen, muscle, testis, placenta and melanocytes, zygin 2 may be involved in the outgrowth and fasciculation of axons. Existing as two alternatively spliced isoforms, the gene encoding zygin 2 maps to human chromosome 2p22.2 and mouse chromosome 17 E3. The genes encoding zygin 1 and 2 are evolutionarily conserved orthologs to the *C. elegans* gene, UNC-76. Zygin 2 is able to restore partial locomotion and axonal fasciculation to *C. elegans* UNC-76 mutants, which suggest a functional role in axonal outgrowth.

## REFERENCES

- Bloom, L. and Horvitz, H.R. 1997. The *Caenorhabditis elegans* gene unc-76 and its human homologs define a new gene family involved in axonal outgrowth and fasciculation. Proc. Natl. Acad. Sci. USA 94: 3414-3419.
- Fujita, T., Ikuta, J., Hamada, J., Okajima, T., Tatematsu, K., Tanizawa, K. and Kuroda, S. 2004. Identification of a tissue-non-specific homologue of axonal fasciculation and elongation protein  $\zeta$ -1. Biochem. Biophys. Res. Commun. 313: 738-744.
- Assmann, E.M., Alborghetti, M.R., Camargo, M.E. and Kobarg, J. 2006. FEZ1 dimerization and interaction with transcription regulatory proteins involves its coiled-coil region. J. Biol. Chem. 281: 9869-9881.
- Alborghetti, M.R., Furlan, A.S., Silva, J.C., Paes Leme, A.F., Torriani, I.C. and Kobarg, J. 2010. Human FEZ1 protein forms a disulfide bond mediated dimer: implications for cargo transport. J. Proteome Res. 9: 4595-4603.
- Alborghetti, M.R., Furlan, A.S. and Kobarg, J. 2011. FEZ2 has acquired additional protein interaction partners relative to FEZ1: functional and evolutionary implications. PLoS ONE 6: e17426.

## CHROMOSOMAL LOCATION

Genetic locus: Fez2 (mouse) mapping to 17 E3.

## PRODUCT

zygin 2 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see zygin 2 shRNA Plasmid (m): sc-155855-SH and zygin 2 shRNA (m) Lentiviral Particles: sc-155855-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

zygin 2 siRNA (m) is recommended for the inhibition of zygin 2 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

zygin 2 (F-3): sc-390111 is recommended as a control antibody for monitoring of zygin 2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor zygin 2 gene expression knockdown using RT-PCR Primer: zygin 2 (m)-PR: sc-155855-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.