

MRCK γ siRNA (m): sc-155919

BACKGROUND

Protein kinases comprise a large group of encoded factors that regulate cellular processes by catalyzing the transfer of a phosphate group to a hydroxyl acceptor in serine, threonine or tyrosine residues. MRCK γ (serine/threonine-protein kinase MRCK γ), also known as CDC42BPG (Cdc42 binding protein kinase γ (DMPK-like)), myotonic dystrophy protein kinase-like α , DMPK2, HSMDPKIN or KAPPA-200, is a 1,551 amino acid cytoplasmic protein belonging to the protein kinase superfamily. Expressed in skeletal muscle and heart, MRCK γ exists as both a homodimer and homotetramer. MRCK γ may function as a downstream effector of Cdc42 in cytoskeletal reorganization, and is known to regulate the phosphorylation of MYPT1 and MYL2, which is required for actomyosin contractility in cell invasion. MRCK γ binds magnesium as a cofactor and strongly associates with GTP-bound Cdc42.

REFERENCES

- Hunter, T. 1995. Protein kinases and phosphatases: the yin and yang of protein phosphorylation and signaling. *Cell* 80: 225-236.
- Leung, T., Chen, X.Q., Tan, I., Manser, E. and Lim, L. 1998. Myotonic dystrophy kinase-related Cdc42-binding kinase acts as a Cdc42 effector in promoting cytoskeletal reorganization. *Mol. Cell. Biol.* 18: 130-140.
- Nakamura, N., Oshiro, N., Fukata, Y., Amano, M., Fukata, M., Kuroda, S., Matsuura, Y., Leung, T., Lim, L. and Kaibuchi, K. 2000. Phosphorylation of ERM proteins at filopodia induced by Cdc42. *Genes Cells* 5: 571-581.
- Ng, Y., Tan, I., Lim, L. and Leung, T. 2004. Expression of the human myotonic dystrophy kinase-related Cdc42-binding kinase γ is regulated by promoter DNA methylation and Sp1 binding. *J. Biol. Chem.* 279: 34156-34164.
- Garcia, P., Ucurum, Z., Bucher, R., Svergun, D.I., Huber, T., Lustig, A., Konarev, P.V., Marino, M. and Mayans, O. 2006. Molecular insights into the self-assembly mechanism of dystrophin myotonic kinase. *FASEB J.* 20: 1142-1151.
- Choi, S.H., Czifra, G., Kedei, N., Lewin, N.E., Lazar, J., Pu, Y., Marquez, V.E. and Blumberg, P.M. 2008. Characterization of the interaction of phorbol esters with the C1 domain of MRCK (myotonic dystrophy kinase-related Cdc42 binding kinase) α/β . *J. Biol. Chem.* 283: 10543-10549.

CHROMOSOMAL LOCATION

Genetic locus: Cdc42bpg (mouse) mapping to 19 A.

PRODUCT

MRCK γ siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MRCK γ shRNA Plasmid (m): sc-155919-SH and MRCK γ shRNA (m) Lentiviral Particles: sc-155919-V as alternate gene silencing products.

For independent verification of MRCK γ (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155919A and sc-155919B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MRCK γ siRNA (m) is recommended for the inhibition of MRCK γ expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MRCK γ gene expression knockdown using RT-PCR Primer: MRCK γ (m)-PR: sc-155919-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.