Ep-CAM (h): 293T Lysate: sc-159303



The Power to Question

BACKGROUND

The epithelial cell adhesion molecule Ep-CAM, which is also designated tumor-associated calcium signal transducer 1 and MK-1, is a monomeric membrane glycoprotein that is expressed in most normal human epithelium and in most carcinomas. The human Ep-CAM gene encodes a 314 amino acid protein that is expressed as 2 forms, a major form and a minor form, which are reduced upon treatment with the amino-glycosylation inhibitor Tunicamycin. Ep-CAM is overexpressed in a variety of carcinomas and is, therefore, a potential target for the visualization and therapy of human solid tumors. Ep-CAM contains an extracellular domain containing two epidermal growth factor-like repeats, followed by a cysteine poor region, which are necessary for the adhesion properties of the molecule.

REFERENCES

- Farr, A., et al. 1991. Epithelial heterogeneity in the murine thymus: a cell surface glycoprotein expressed by subcapsular and medullary epithelium. J. Histochem. Cytochem. 39: 645-653.
- Bergsagel, P.L., et al. 1992. A murine cDNA encodes a pan-epithelial glycoprotein that is also expressed on plasma cells. J. Immunol. 148: 590-596.
- Bjork, P., et al. 1993. Isolation, partial characterization, and molecular cloning of a human colon adenocarcinoma cell-surface glycoprotein recognized by the C215 mouse monoclonal antibody. J. Biol. Chem. 268: 24232-24241.
- 4. Nelson, A.J., et al. 1996. The murine homolog of human Ep-CAM, a homotypic adhesion molecule, is expressed by thymocytes and thymic epithelial cells. Eur. J. Immunol. 26: 401-408.
- Litvinov, S.V., et al. 1997. Epithelial cell adhesion molecule (Ep-CAM) modulates cell-cell interactions mediated by classic cadherins. J. Cell Biol. 139: 1337-1348.
- Tomita, Y., et al. 2000. Molecular identification of a human carcinomaassociated glycoprotein antigen recognized by mouse monoclonal antibody FU-MK-1. J. Cancer Res. 91: 231-238.
- 7. Taguchi, N., et al. 2000. Abnormal thymic expression of epithelial cell adhesion molecule (Ep-CAM) in New Zealand Black (NZB) mice. J. Autoimmun. 13: 393-404.
- Trebak, M., et al. 2001. Oligomeric state of the colon carcinoma-associated glycoprotein GA733-2 (Ep-CAM/EGP40) and its role in GA733-mediated homotypic cell-cell adhesion. J. Biol. Chem. 276: 2299-2309.
- Balzar, M., et al. 2001. Epidermal growth factor-like repeats mediate lateral and reciprocal interactions of Ep-CAM molecules in homophilic adhesions. Mol. Cell. Biol. 21: 2570-2580.

CHROMOSOMAL LOCATION

Genetic locus: EPCAM (human) mapping to 2p21.

PRODUCT

Ep-CAM (h): 293T Lysate represents a lysate of human Ep-CAM transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Ep-CAM (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Ep-CAM antibodies. Recommended use: 10-20 μ l per lane


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Ep-CAM (C-10): sc-25308 is recommended as a positive control antibody for Western Blot analysis of enhanced human Ep-CAM expression in Ep-CAM transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG κ BP-HRP: sc-516102 or m-lgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Ep-CAM (C-10): sc-25308. Western blot analysis of Ep-CAM expression in non-transfected: sc-117752 (A) and human Ep-CAM transfected: sc-159303 (B) 293T whole cell lyestes

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com