



Bmx (h2): 293T Lysate: sc-159692

BACKGROUND

The Tec family of non-receptor tyrosine kinases is composed of six proteins, designated Tec, Emt (also known as Itk or Tsk), Btk (previously known as Atk, BPK or Emb), Bmx, Txk (also known as Rlk) and Dsrc28C. All members of the family contain SH3 and SH2 domains and, with the exception of Txk and Dsrc28C, also contain a Pleckstrin homology (PH) and a Tec homology (TH) domain in their amino-termini. Four alternatively spliced forms of Tec are found to be expressed broadly in cells of hematopoietic lineage and in hepatocytes. The Emt gene product associates with CD28 and becomes activated subsequent to CD28 ligation. Btk is necessary for proper B cell development, and mutations in the gene encoding Btk have been associated with families suffering from X-linked agammaglobulinemia, also referred to as Bruton's disease. The Bmx protein shares a high degree of homology with Btk and seems to be expressed at highest levels in the heart. Txk expression is T cell-specific, while expression of the *Drosophila* Tec homolog, Dsrc28C, is developmentally regulated.

REFERENCES

1. Yamada, N., et al. 1993. Structure and expression of novel protein tyrosine kinases, Emb and Emt, in hematopoietic cells. *Biochem. Biophys. Res. Commun.* 192: 231-240.
2. Thomas, J.D., et al. 1993. Colocalization of X-linked agammaglobulinemia and X-linked immunodeficiency genes. *Science* 261: 355-358.
3. Tamagnone, L., et al. 1994. Bmx, a novel nonreceptor tyrosine kinase gene of the Btk/Itk/Tec/Txk family located in chromosome Xp22.2. *Oncogene* 9: 3683-3688.
4. Haire, R.N., et al. 1994. Txk, a novel human tyrosine kinase expressed in T cells shares sequence identity with Tec family kinases and maps to 4p12. *Hum. Mol. Genet.* 3: 897-901.
5. August, A., et al. 1994. CD28 is associated with and induces the immediate tyrosine phosphorylation and activation of the Tec family kinase Itk/Emt in the human Jurkat leukemic T-cell line. *Proc. Natl. Acad. Sci. USA* 91: 9347-9351.
6. Hu, Q., et al. 1995. Identification of Rlk, a novel protein tyrosine kinase with predominant expression in the T cell lineage. *J. Biol. Chem.* 270: 1928-1934.

CHROMOSOMAL LOCATION

Genetic locus: BMX (human) mapping to Xp22.2.

PRODUCT

Bmx (h2): 293T Lysate represents a lysate of human Bmx transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

Bmx (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Bmx antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.