

PAOX (h2): 293T Lysate: sc-159701

BACKGROUND

PAOX (polyamine oxidase (exo-N₄-amino)), also known as PAO, is a 649 amino acid protein that localizes to both the cytoplasm and to peroxisomes and belongs to the flavin monoamine oxidase family. Expressed in a variety of tissues, PAOX exists as a monomer that uses FAD as a cofactor to catalyze the oxidation of N₁-acetylspermine and N₁-acetylspermidine to spermidine and putrescine, respectively, thereby playing a role in polyamine back-conversion, as well as in the regulation of polyamine intracellular concentration. Multiple isoforms of PAOX exist due to alternative splicing events. The gene encoding PAOX maps to human chromosome 10, which houses over 1,200 genes and comprises nearly 4.5% of the human genome. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromic deafness, Wolman's syndrome, Cowden syndrome, multiple endocrine neoplasia type 2 and porphyria.

REFERENCES

1. Seiler, N. 1995. Polyamine oxidase, properties and functions. *Prog. Brain Res.* 106: 333-344.
2. Vujcic, S., Diegelman, P., Bacchi, C.J., Kramer, D.L. and Porter, C.W. 2002. Identification and characterization of a novel flavin-containing spermine oxidase of mammalian cell origin. *Biochem. J.* 367: 665-675.
3. Vujcic, S., Liang, P., Diegelman, P., Kramer, D.L. and Porter, C.W. 2003. Genomic identification and biochemical characterization of the mammalian polyamine oxidase involved in polyamine back-conversion. *Biochem. J.* 370: 19-28.
4. Casero, R.A., Wang, Y., Stewart, T.M., Devereux, W., Hacker, A., Wang, Y., Smith, R. and Woster, P.M. 2003. The role of polyamine catabolism in anti-tumour drug response. *Biochem. Soc. Trans.* 31: 361-365.
5. Wu, T., Yankovskaya, V. and McIntire, W.S. 2003. Cloning, sequencing, and heterologous expression of the murine peroxisomal flavoprotein, N₁-acetylated polyamine oxidase. *J. Biol. Chem.* 278: 20514-20525.
6. Pledgie, A., Huang, Y., Hacker, A., Zhang, Z., Woster, P.M., Davidson, N.E. and Casero, R.A. 2005. Spermine oxidase SMO(PAOh1), Not N₁-acetylpolyamine oxidase PAO, is the primary source of cytotoxic H₂O₂ in polyamine analogue-treated human breast cancer cell lines. *J. Biol. Chem.* 280: 39843-39851.
7. Järvinen, A., Keinänen, T.A., Grigorenko, N.A., Khomutov, A.R., Uimari, A., Vepsäläinen, J., Närvänen, A., Alhonen, L. and Jänne, J. 2006. Guide molecule-driven stereospecific degradation of α -methylpolyamines by polyamine oxidase. *J. Biol. Chem.* 281: 4589-4595.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: PAOX (human) mapping to 10q26.3.

PRODUCT

PAOX (h2): 293T Lysate represents a lysate of human PAOX transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

PAOX (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive PAOX antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.