

AGTRAP (h): 293T Lysate: sc-159795

BACKGROUND

AGTRAP (Angiotensin II receptor-associated protein), also known as ATRAP, is a transmembrane protein that localizes to the Golgi apparatus, the endoplasmic reticulum (ER), endocytotic vesicles and perinuclear vesicular structures. Highly expressed in heart, kidney, pancreas and thyroid, AGTRAP functions as a negative regulator of the Angiotensin II type I receptor (AT1). AGTRAP controls receptor internalization and receptor desensitization events (such as phosphorylation) and, through this control, decreases Angiotensin II signaling, thereby reducing rates of cell proliferation and Angiotensin II-stimulated transcriptional activity. AGTRAP is 159 amino acids in length and is able to bind RACK1 (receptor for activated C kinase 1); an association that is thought to help recruit AGTRAP to AT1. Two isoforms of AGTRAP exist due to alternative splicing events.

REFERENCES

1. Daviet, L., et al. 1999. Cloning and characterization of ATRAP, a novel protein that interacts with the Angiotensin II type 1 receptor. *J. Biol. Chem.* 274: 17058-17062.
2. Cui, T., et al. 2000. ATRAP, novel AT1 receptor associated protein, enhances internalization of AT1 receptor and inhibits vascular smooth muscle cell growth. *Biochem. Biophys. Res. Commun.* 279: 938-941.
3. Wang, W., et al. 2002. Identification and characterization of AGTRAP, a human homolog of murine Angiotensin II receptor-associated protein (AGTRAP). *Int. J. Biochem. Cell Biol.* 34: 93-102.
4. Lopez-Illasaca, M., et al. 2003. The Angiotensin II type I receptor-associated protein, ATRAP, is a transmembrane protein and a modulator of Angiotensin II signaling. *Mol. Biol. Cell* 14: 5038-5050.
5. Guo, D.F., et al. 2003. Type 1 Angiotensin II receptor-associated protein ARAP1 binds and recycles the receptor to the plasma membrane. *Biochem. Biophys. Res. Commun.* 310: 1254-1265.
6. Guo, D.F., et al. 2004. A novel angiotensin II type 1 receptor-associated protein induces cellular hypertrophy in rat vascular smooth muscle and renal proximal tubular cells. *J. Biol. Chem.* 279: 21109-21120.
7. Guo, S., et al. 2005. Identification of calcium-modulating cyclophilin ligand (CAML) as transducer of angiotensin II-mediated nuclear factor of activated T cells (NFAT) activation. *J. Biol. Chem.* 280: 12536-12541.
8. Tanaka, Y., et al. 2005. The novel angiotensin II type 1 receptor (AT1R)-associated protein ATRAP downregulates AT1R and ameliorates cardiomyocyte hypertrophy. *FEBS Lett.* 579: 1579-1586.

CHROMOSOMAL LOCATION

Genetic locus: AGTRAP (human) mapping to 1p36.22.

PRODUCT

AGTRAP (h): 293T Lysate represents a lysate of human AGTRAP transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

AGTRAP (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive AGTRAP antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.