

Rent1 (E-8): sc-166091

BACKGROUND

In eukaryotes, it is essential to have the ability to detect and degrade transcripts that lack full coding potential. Nonsense-mediated RNA decay (NMD) protects the organism by avoiding the translation of truncated peptides with dominant negative or deleterious gain-of-function potential. Rent1, a mammalian ortholog of Upf1, is essential for embryonic viability. Rent1 (also designated regulator of nonsense transcripts and HUPf1) contains an N-terminal zinc finger-like domain, NTPase domains and a region comprised of domains that define Rent1 as a superfamily group I helicase. Rent1 protein has nucleic-acid-dependent ATPase activity and 5' to 3' helicase activity. In addition, Rent1 is an RNA-binding protein whose activity is modulated by ATP and directly interacts with Rent2, which is a mammalian homolog of Upf2p. Two mammalian orthologs to Upf3p, Rent3a and Rent3b, are encoded by two separate genes. Rent3b (also known as Rent3X) is encoded by a X-linked gene and localizes primarily to the nucleus, while Rent 1 and Rent 2 localize primarily in the cytoplasm. Specific Rent3 protein interactions with Y14 and spliced mRNA suggest Rent3a and Rent3b serve as a link between splicing and NMD in the cytoplasm.

CHROMOSOMAL LOCATION

Genetic locus: UPF1 (human) mapping to 19p13.11; Upf1 (mouse) mapping to 8 B3.3.

SOURCE

Rent1 (E-8) is a mouse monoclonal antibody raised against amino acids 71-370 mapping near the N-terminus of Rent1 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-166091 X, 200 µg/0.1 ml.

APPLICATIONS

Rent1 (E-8) is recommended for detection of Rent1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Rent1 siRNA (h): sc-38223, Rent1 siRNA (m): sc-38224, Rent1 shRNA Plasmid (h): sc-38223-SH, Rent1 shRNA Plasmid (m): sc-38224-SH, Rent1 shRNA (h) Lentiviral Particles: sc-38223-V and Rent1 shRNA (m) Lentiviral Particles: sc-38224-V.

Rent1 (E-8) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of Rent1: 130 kDa.

Positive Controls: Sol8 cell lysate: sc-2249, A-673 cell lysate: sc-2414 or C2C12 whole cell lysate: sc-364188.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



Rent1 (E-8): sc-166091. Western blot analysis of Rent1 expression in Sol8 (A), C2C12 (B), A-673 (C), Ramos (D) and NAMALWA (E) whole cell lysates.



Rent1 (E-8): sc-166091. Immunoperoxidase staining of formalin fixed, paraffin-embedded human brain tissue showing cytoplasmic staining of neuronal cells.

SELECT PRODUCT CITATIONS

1. Wang, G., et al. 2016. MiR-128 and miR-125 regulate expression of coagulation Factor IX gene with nonsense mutation by repressing nonsense-mediated mRNA decay. *Biomed. Pharmacother* 80: 331-337.
2. Raguz, J., et al. 2016. Epidermal RAF prevents allergic skin disease. *Elife* 5: e14012.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.