

HoxD10 (G-3): sc-166235

BACKGROUND

The Hox proteins play a role in development and cellular differentiation by regulating downstream target genes. Specifically, the Hox proteins direct DNA-protein and protein-protein interactions that assist in determining the morphologic features associated with the anterior-posterior body axis. Hox proteins are involved in controlling axial patterning, leukaemias and hereditary malformations. Homeobox protein HoxD10, also designated Hox-4D or Hox-4E, belongs to the Abd-B homeobox family of proteins. HoxD10 is a nuclear protein primarily expressed in the adult male and female urogenital tracts but also expressed in developing limb buds during development. Defects in the gene encoding for the HoxD10 protein cause congenital vertical talus (CVT), more commonly known as rocker-bottom foot deformity. CVT is characterized by a dislocation of the talonavicular joint but is usually accompanied by other congenital deformities.

REFERENCES

1. Zappavigna, V., et al. 1991. Hox4 genes encode transcription factors with potential auto- and cross-regulatory capacities. *EMBO J.* 10: 4177-4187.
2. Redline, R.W., et al. 1992. Human Hox4E: a gene strongly expressed in the adult male and female urogenital tracts. *Genomics* 13: 425-430.
3. Gabellini, D., et al. 2003. Early mitotic degradation of the homeoprotein HoxC10 is potentially linked to cell cycle progression. *EMBO J.* 22: 3715-3724.
4. Juan, A.H., et al. 2003. Enhancer timing of Hox gene expression: deletion of the endogenous HoxC8 early enhancer. *Development* 130: 4823-4834.
5. Miller, G.J., et al. 2003. Aberrant HoxC expression accompanies the malignant phenotype in human prostate. *Cancer Res.* 63: 5879-5888.

CHROMOSOMAL LOCATION

Genetic locus: HOXD10 (human) mapping to 2q31.1; Hoxd10 (mouse) mapping to 2 C3.

SOURCE

HoxD10 (G-3) is a mouse monoclonal antibody raised against amino acids 21-100 mapping near the N-terminus of HoxD10 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-166235 X, 200 µg/0.1 ml.

HoxD10 (G-3) is available conjugated to agarose (sc-166235 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-166235 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-166235 PE), fluorescein (sc-166235 FITC), Alexa Fluor® 488 (sc-166235 AF488), Alexa Fluor® 546 (sc-166235 AF546), Alexa Fluor® 594 (sc-166235 AF594) or Alexa Fluor® 647 (sc-166235 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-166235 AF680) or Alexa Fluor® 790 (sc-166235 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

HoxD10 (G-3) is recommended for detection of HoxD10 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HoxD10 (G-3) is also recommended for detection of HoxD10 in additional species, including equine, canine, bovine and porcine.


Suitable for use as control antibody for HoxD10 siRNA (h): sc-44814, HoxD10 siRNA (m): sc-44815, HoxD10 shRNA Plasmid (h): sc-44814-SH, HoxD10 shRNA Plasmid (m): sc-44815-SH, HoxD10 shRNA (h) Lentiviral Particles: sc-44814-V and HoxD10 shRNA (m) Lentiviral Particles: sc-44815-V.

HoxD10 (G-3) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of HoxD10: 40 kDa.

Positive Controls: HoxD10 (m2): 293T Lysate: sc-126972 or ECV304 cell lysate: sc-2269.

DATA



HoxD10 (G-3): sc-166235. Western blot analysis of HoxD10 expression in non-transfected: sc-117752 (A) and mouse HoxD10 transfected: sc-126972 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

1. Liang, A.L., et al. 2016. miRNA-10b sponge: an anti-breast cancer study *in vitro*. *Oncol. Rep.* 35: 1950-1958.
2. Tupone, M.G., et al. 2020. microRNA-378a-5p is a novel positive regulator of melanoma progression. *Oncogenesis* 9: 22.
3. Yu, M., et al. 2024. Hsa_circ_0000825 promotes the progression of laryngeal squamous cell carcinoma by sponging miR-766 and interacting with ELAVL1. *Heliyon* 10: e37264.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.