SANTA CRUZ BIOTECHNOLOGY, INC.

Dvl (B-4): sc-166303



BACKGROUND

Mammalian homologs of the *Drosophila* dishevelled (Dsh) gene have been identified, including DvI-1, DvI-2 and DvI-3. The mammalian dishevelled proteins contain three homologous domains, two of which are unrelated to any other known protein. The third region is homologous to the discs-large homology domain of *Drosophila* discs-large-1, a tumor suppressor protein. Like their *Drosophila* counterpart, the dishevelled proteins are thought to be involved in embryogenesis. Overexpression of DvI-1 has been shown to inhibit the phosphorylation of Tau by GSK-3 β . This finding may prove to be important in Alzheimer's studies, which have shown that Tau is hyperphosphorylated. In *Drosophila*, Dsh is a component of the frizzled signaling pathway.

SOURCE

Dvl (B-4) is a mouse monoclonal antibody raised against amino acids 21-280 mapping near the N-terminus of Dvl-1 of human origin.

PRODUCT

Each vial contains 200 μg IgG_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Dvl (B-4) is available conjugated to agarose (sc-166303 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-166303 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-166303 PE), fluorescein (sc-166303 FITC), Alexa Fluor® 488 (sc-166303 AF488), Alexa Fluor® 546 (sc-166303 AF546), Alexa Fluor® 594 (sc-166303 AF594) or Alexa Fluor® 647 (sc-166303 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-166303 AF680) or Alexa Fluor® 790 (sc-166303 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

Dvl (B-4) is recommended for detection of Dvl-1, Dvl-2 and Dvl-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Dvl: 85 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG K BP-HRP: sc-516102 or m-IgG K BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG K BP-FITC: sc-516140 or m-IgG K BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA





Dvl (B-4) HRP: sc-166303 HRP. Direct western blot analysis of Dvl expression in F9 (A), HeLa (B), MDA-MB-231 (C), KNRK (D) and BT-20 (E) whole cell lysates.

Dvl (B-4): sc-166303. Western blot analysis of Dvl expression in MDA-MB-231 (A), Heta (B) and PC-3 (C) whole cell lysates and rat testis (D) and human skeletal muscle (E) tissue extracts. Detection reagent used: m-lgG₁ BP-HRP: sc-525408.

SELECT PRODUCT CITATIONS

- 1. Arellanes-Robledo, J., et al. 2013. Fibrogenic actions of acetaldehyde are β -catenin dependent but Wingless independent: a critical role of nucleoredoxin and reactive oxygen species in human hepatic stellate cells. Free Radic. Biol. Med. 65: 1487-1496.
- Shimizu, N., et al. 2014. Hipk2 and PP1c cooperate to maintain Dvl protein levels required for Wnt signal transduction. Cell Rep. 8: 1391-1404.
- Yeh, C.C., et al. 2015. Genistein suppresses growth of human uterine sarcoma cell lines via multiple mechanisms. Anticancer Res. 35: 3167-3173.
- Zhang, Q., et al. 2017. The preventive effects of taurine on neural tube defects through the Wnt/PCP-Jnk-dependent pathway. Amino Acids 49: 1633-1640.
- Aznar, N., et al. 2018. Convergence of Wnt, growth factor, and heterotrimeric G protein signals on the guanine nucleotide exchange factor DAPLE. Sci. Signal. 11: eaao4220.
- Yang, Y., et al. 2019. Endogenous IGF signaling directs heterogeneous mesoderm differentiation in human embryonic stem cells. Cell Rep. 29: 3374-3384.e5.
- Kawarazaki, W., et al. 2020. Salt causes aging-associated hypertension via vascular Wnt5a under Klotho deficiency. J. Clin. Invest. 130: 4152-4166.
- He, W., et al. 2021. Activation of Frizzled-7 attenuates blood-brain barrier disruption through Dvl/β-catenin/WISP1 signaling pathway after intracerebral hemorrhage in mice. Fluids Barriers CNS 18: 44.
- Carr, H.S., et al. 2023. The Wnt pathway protein Dvl1 targets somatostatin receptor 2 for lysosome-dependent degradation. J. Biol. Chem. 299: 104645.

RESEARCH USE

For research use only, not for use in diagnostic procedures.