

FcRH3 (A-12): sc-166347

BACKGROUND

The Fc receptor homolog (FcRH) family of proteins are related to the classical Fc receptors (FcR) and belong to the immunoglobulin receptor superfamily. The proteins in the FcRH family are type I transmembrane glycoproteins and consist of FcRH1-FcRH6. The genes encoding the human FcRH proteins map to chromosome 1, near the related FCR genes. The FcRH proteins, which are involved in immune system regulation, have immunoreceptor-tyrosine inhibitory motifs in their cytoplasmic domains. Mutations in the gene encoding for the FcRH proteins may be associated with systemic lupus erythematosus, autoimmune thyroid disease and rheumatoid arthritis. The FcRH genes are expressed primarily, although not exclusively, by mature B lineage cells, and may serve important regulatory roles in normal and neoplastic B cell development.

REFERENCES

1. Davis, R.S., Wang, Y.H., Kubagawa, H. and Cooper, M.D. 2001. Identification of a family of Fc receptor homologs with preferential B cell expression. *Proc. Natl. Acad. Sci. USA* 98: 9772-9777.
2. Davis, R.S., Dennis, G., Jr., Odom, M.R., Gibson, A.W., Kimberly, R.P., Burrows, P.D. and Cooper, M.D. 2002. Fc receptor homologs: newest members of a remarkably diverse Fc receptor gene family. *Immunol. Rev.* 190: 123-136.
3. Ehrhardt, G.R., Davis, R.S., Hsu, J.T., Leu, C.M., Ehrhardt, A. and Cooper, M.D. 2003. The inhibitory potential of Fc receptor homolog 4 on memory B cells. *Proc. Natl. Acad. Sci. USA* 100: 13489-13494.
4. Davis, R.S., Stephan, R.P., Chen, C.C., Dennis, G., Jr. and Cooper, M.D. 2004. Differential B cell expression of mouse Fc receptor homologs. *Int. Immunol.* 16: 1343-1353.
5. Davis, R.S., Ehrhardt, G.R., Leu, C.M., Hirano, M. and Cooper, M.D. 2005. An extended family of Fc receptor relatives. *Eur. J. Immunol.* 35: 674-680.

CHROMOSOMAL LOCATION

Genetic locus: FCRL3 (human) mapping to 1q23.1.

SOURCE

FcRH3 (A-12) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 631-660 near the C-terminus of FcRH3 of human origin.

PRODUCT

Each vial contains 200 µg IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-166347 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

FcRH3 (A-12) is recommended for detection of FcRH3 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FcRH3 siRNA (h): sc-45791, FcRH3 shRNA Plasmid (h): sc-45791-SH and FcRH3 shRNA (h) Lentiviral Particles: sc-45791-V.


Molecular Weight of FcRH3: 92 kDa.

Positive Controls: Raji whole cell lysate, THP-1 cell lysate: sc-2238 or U-937 cell lysate: sc-2239.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



FcRH3 (A-12): sc-166347. Western blot analysis of FcRH3 expression in THP-1 whole cell lysate.



FcRH3 (A-12): sc-166347. Immunofluorescence staining of methanol-fixed HeLa cells showing membrane localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.