

CA XIV (A-4): sc-166438



The Power to Question

BACKGROUND

Carbonic anhydrases (CAs) are members of a large family of zinc metallo-enzymes that catalyze the reversible hydration of carbon dioxide. CAs are involved in a variety of biological processes, including respiration, calcification, acid-base balance and bone resorption, as well as the formation of aqueous humor, cerebrospinal fluid, saliva and gastric juice. They show extensive diversity in distribution and in their subcellular localization. The human CA2 gene, which maps to chromosome 8q22, encodes CA II, a cytoplasmic protein that has the highest turnover rate and widest tissue distribution of any known human CA isozyme. The human CA4 gene, which maps to chromosome 17q23, encodes CA IV, a membrane-anchored isozyme that is expressed on the luminal surfaces of pulmonary capillaries and proximal renal tubules. The human CA9, CA12 and CA14 genes, which map to chromosomes 9p13.3, 15q22 and 1q21.2, respectively, encode transmembrane proteins that have unique patterns of tissue-specific expression. CA IX is specifically expressed in clear-cell renal carcinomas, whereas CA XII is highly expressed in normal tissues, such as kidney, colon and pancreas. Human CA XIV is also expressed in normal tissues, such as brain, but differs from CA XII in its expression pattern.

REFERENCES

1. Dodgson, S.J., et al., eds. 1991. The Carbonic Anhydrases: Cellular Physiology and Molecular Genetics. New York: Plenum Publishing Corporation.
2. Venta, P.J., et al. 1991. Carbonic anhydrase II deficiency syndrome in a Belgian family is caused by a point mutation at an invariant histidine residue (107 His—Tyr): complete structure of the normal human CA II gene. *Am. J. Hum. Genet.* 49: 1082-1090.

CHROMOSOMAL LOCATION

Genetic locus: CA14 (human) mapping to 1q21.2; Car14 (mouse) mapping to 3 F2.1.

SOURCE

CA XIV (A-4) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 21-48 near the N-terminus of CA XIV of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CA XIV (A-4) is available conjugated to agarose (sc-166438 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-166438 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-166438 PE), fluorescein (sc-166438 FITC), Alexa Fluor® 488 (sc-166438 AF488), Alexa Fluor® 546 (sc-166438 AF546), Alexa Fluor® 594 (sc-166438 AF594) or Alexa Fluor® 647 (sc-166438 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-166438 AF680) or Alexa Fluor® 790 (sc-166438 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-166438 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

CA XIV (A-4) is recommended for detection of CA XIV of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CA XIV (A-4) is also recommended for detection of CA XIV in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for CA XIV siRNA (h): sc-41465, CA XIV siRNA (m): sc-41466, CA XIV shRNA Plasmid (h): sc-41465-SH, CA XIV shRNA Plasmid (m): sc-41466-SH, CA XIV shRNA (h) Lentiviral Particles: sc-41465-V and CA XIV shRNA (m) Lentiviral Particles: sc-41466-V.

Molecular Weight of CA XIV: 50 kDa.


Positive Controls: A2058 whole cell lysate: sc-364178 or AN3 CA cell lysate: sc-24662.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



CA XIV (A-4): sc-166438. Western blot analysis of CA XIV expression in A2058 (A) and AN3 CA (B) whole cell lysates. Detection reagent used: m-IgGκ BP-HRP: sc-516102.



CA XIV (A-4): sc-166438. Near-infrared western blot analysis of CA XIV expression in A2058 (A) and AN3 CA (B) whole cell lysates. Blocked with UltraCruz® Blocking Reagent: sc-516214. Detection reagent used: m-IgGκ BP-CFL 680: sc-516180.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA