

SSX (C-7): sc-166595

BACKGROUND

The transcriptional coactivator SYT (synovial translocation protein) contains a conserved amino terminal SNH domain and a carboxy terminal QPGY domain, which is a functioning transcriptional activating sequence. Synovial sarcoma translocation (SSX) proteins, including SSX1-5, are transcriptional repressors that contain a repressor domain in their carboxy termini. SSX proteins are localized to the nucleus and expressed in testis and several types of cancers and, therefore, they are classified as C/T (cancer/testis) antigens. The t(x;18) translocation results in the fusion of the amino terminus of SYT to the carboxy terminus of either SSX1 or SSX2; both fusions result in the production of transcriptional activators. SYT-SSX chimeras are detected in most synovial sarcomas. Synovial sarcomas are responsible for up to 10% of soft tissue sarcomas and are histologically characterized as either biphasic or monophasic. Genetic analysis indicates that biphasic synovial sarcomas contain SYT-SSX1 fusions, whereas SYT-SSX2 fusions are found in monophasic synovial sarcomas, providing additional distinguishing characterization of these subtypes.

REFERENCES

- Clark, J., et al. 1994. Identification of novel genes, SYT and SSX, involved in the t(X;18)(p11.2;q11.2) translocation found in human synovial sarcoma. *Nat. Genet.* 7: 502-508.
- Crew, A.J., et al. 1995. Fusion of SYT to two genes, SSX1 and SSX2, encoding proteins with homology to the Krüppel-associated box in human synovial sarcoma. *EMBO J.* 14: 2333-2340.
- dos Santos, N.R., et al. 1997. Nuclear localization of SYT, SSX and the synovial sarcoma-associated SYT-SSX fusion proteins. *Hum. Mol. Genet.* 6: 1549-1558.

CHROMOSOMAL LOCATION

Genetic locus: SSX1/SSX3/SSX4/SSX4B/SSX5/SSX6/SSX9 (human) mapping to Xp11.23, SSX2/SSX2B/SSX7/SSX8 (human) mapping to Xp11.22.

SOURCE

SSX (C-7) is a mouse monoclonal antibody raised against amino acids 1-188 representing full length SSX1 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-166595 X, 200 µg/0.1 ml.

SSX (C-7) is available conjugated to agarose (sc-166595 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-166595 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-166595 PE), fluorescein (sc-166595 FITC), Alexa Fluor® 488 (sc-166595 AF488), Alexa Fluor® 546 (sc-166595 AF546), Alexa Fluor® 594 (sc-166595 AF594) or Alexa Fluor® 647 (sc-166595 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-166595 AF680) or Alexa Fluor® 790 (sc-166595 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

SSX (C-7) is recommended for detection of SSX1-9 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SSX (C-7) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of SSX: 22 kDa.

Positive Controls: TT whole cell lysate: sc-364195, NCI-H929 whole cell lysate: sc-364786 or SSX (h2): 293T Lysate: sc-112491.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



SSX (C-7): sc-166595. Western blot analysis of SSX expression in non-transfected 293T: sc-117752 (A), human SSX transfected 293T: sc-112491 (B) and TT (C) whole cell lysates.



SSX (C-7): sc-166595. Western blot analysis of SSX expression in NCI-H929 whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.