

ERp57 (B-5): sc-166680

BACKGROUND

Mammals defend themselves against intracellular pathogens through presentation of cytoplasmically-derived short pathogenic peptides to the cell surface of cytotoxic T lymphocytes, which subsequently leads to cytotoxic events with respect to the affected cell. Antigen presentation is mediated by major histocompatibility complex (MHC) class I molecules, which bind and coordinate short pathogenic peptides. MHC class I molecules assemble in the endoplasmic reticulum with chaperones before binding to the transporter associated with antigen processing (TAP). ERp57, also designated GRP57, GRP58, ERp60 and ERp61, is a component of the MHC class I pathway that appears to interact with MHC class I molecules before they associate with TAP. The human ERp57 gene maps to chromosome 15q15.3 and encodes a 505 amino acid protein. ERp57 has two Trp-Cys-Gly-His-Cys-Lys motifs completely conserved among the mammals. ERp57 may act as a protease, a protein disulfide isomerase, a phospholipase or a combination of these.

CHROMOSOMAL LOCATION

Genetic locus: PDIA3 (human) mapping to 15q15.3; Pdia3 (mouse) mapping to 2 E5.

SOURCE

ERp57 (B-5) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 20-60 near the N-terminus of ERp57 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-166680 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

ERp57 (B-5) is recommended for detection of ERp57 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ERp57 (B-5) is also recommended for detection of ERp57 in additional species, including canine.

Suitable for use as control antibody for ERp57 siRNA (h): sc-35341, ERp57 siRNA (m): sc-42876, ERp57 siRNA (r): sc-270455, ERp57 shRNA Plasmid (h): sc-35341-SH, ERp57 shRNA Plasmid (m): sc-42876-SH, ERp57 shRNA Plasmid (r): sc-270455-SH, ERp57 shRNA (h) Lentiviral Particles: sc-35341-V, ERp57 shRNA (m) Lentiviral Particles: sc-42876-V and ERp57 shRNA (r) Lentiviral Particles: sc-270455-V.


Molecular Weight of ERp57: 61 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Daudi cell lysate: sc-2415 or Caki-1 cell lysate: sc-2224.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



ERp57 (B-5): sc-166680. Western blot analysis of ERp57 expression in HeLa (A), Daudi (B), Caki-1 (C), KNRK (D) and Hep G2 (E) whole cell lysates.



ERp57 (B-5): sc-166680. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human thyroid gland tissue showing cytoplasmic staining of glandular cells (B).

SELECT PRODUCT CITATIONS

1. Rocchiccioli, S., et al. 2012. Proteomics changes in adhesion molecules: a driving force for vascular smooth muscle cell phenotypic switch. *Mol. Biosyst.* 8: 1052-1059.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.