

MASP-1/3 (G-7): sc-166816

BACKGROUND

Mannose (or mannan)-binding lectin (MBL), also known as serum mannose-binding protein (MBP), initiates the lectin branch of the innate immune response by binding to the surface of potentially pathogenic microorganisms and initiating complement fixation through an N-terminal collagen-like domain. MBL is a key component in immune response in that it can directly trigger neutralization of invading microorganisms by an Ab-independent mechanism. Mutations of human MBL are associated with immunodeficiency resulting from a reduction in the ability of the mutant MBL to initiate complement fixation. In human, three types of MBL-associated serine proteases, MASP-1, MASP-2 and MASP-3, and a truncated form of MASP-2 (small MBL-associated protein; sMAP or Map19) complex with MBL to activate the lectin pathway of the complement system. MASP-3 is an alternatively spliced product from the MASP-1 gene. The heavy/A chains are identical between MASP-1 and MASP-3 but the light/B chains are entirely different. Activated MASPs subsequently cleave and activate downstream components of the complement pathway.

REFERENCES

1. Heise, C., et al. 2000. Impaired secretion of rat mannose-binding protein resulting from mutations in the collagen-like domain. *J. Immunol.* 165: 1403-1409.
2. Matsushita, M., et al. 2000. Proteolytic activities of two types of mannose-binding lectin-associated serine protease. *J. Immunol.* 165: 2637-2642.
3. Chen, C.B. and Wallis, R. 2001. Stoichiometry of complexes between mannose-binding protein and its associated serine proteases: defining functional units for complement activation. *J. Biol. Chem.* 276: 25894-25902.
4. Endo, M., et al. 2001. Regulation of *in situ* complement activation via the lectin pathway in patients with IgA nephropathy. *Clin. Nephrol.* 55: 185-191.
5. Thielens, N.M., et al. 2001. Interaction properties of human mannan-binding lectin (MBL)-associated serine proteases-1 and -2, MBL-associated protein 19, and MBL. *J. Immunol.* 166: 5068-5077.
6. Dahl, M.R., et al. 2001. MASP-3 and its association with distinct complexes of the mannan-binding lectin complement activation pathway. *Immunity* 15: 127-135.

CHROMOSOMAL LOCATION

Genetic locus: MASP1 (human) mapping to 3q27.3.

SOURCE

MASP-1/3 (G-7) is a mouse monoclonal antibody raised against amino acids 171-430 mapping within an internal region of MASP-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

MASP-1/3 (G-7) is recommended for detection of MASP-1/3 heavy chain of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MASP-1/3 siRNA (h): sc-45349, MASP-1/3 shRNA Plasmid (h): sc-45349-SH and MASP-1/3 shRNA (h) Lentiviral Particles: sc-45349-V.

Molecular Weight of MASP-1/3 proenzyme: 90 kDa.

Molecular Weight of MASP-1/3 heavy chain: 65 kDa.


Molecular Weight of MASP-1/3 light chain: 36 kDa.

Positive Controls: BT-20 cell lysate: sc-2223.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



MASP-1/3 (G-7): sc-166816. Western blot analysis of MASP-1/3 expression in BT-20 whole cell lysate.



MASP-1/3 (G-7): sc-166816. Western blot analysis of full-length human recombinant MASP-1.

SELECT PRODUCT CITATIONS

1. Trudel, D., et al. 2019. Proteases and their inhibitors as prognostic factors for high-grade serous ovarian cancer. *Pathol. Res. Pract.* 215: 152369.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.