

# p-c-Src (H-8): sc-166859

## BACKGROUND

The major translational products of the Src gene family are membrane-associated tyrosine protein kinases that lack transmembrane and external amino acid sequences. By virtue of their common structural motifs, the Src family is composed of nine members in vertebrates, including c-Src, c-Yes, Fgr, Yrk, Fyn, Lyn, Hck, Lck and Blk. Src family kinases, which contain an amino-terminal cell membrane anchor followed by SH3 and SH2 domains, transduce signals that are involved in the control of a variety of cellular processes, including proliferation, differentiation, motility and adhesion. Src family members are normally maintained in an inactive state and can be activated transiently during cellular events such as mitosis. Different subcellular locations of Src family kinases may be important for the regulation of specific cellular processes, such as mitogenesis, cytoskeletal organization and membrane trafficking. c-Src (also designated pp60Src, Src p60 and proto-oncogene tyrosine protein kinase Src) is expressed in a broad range of tissue and cell types, although the highest levels of c-Src are detected in neuronal tissues and platelets. c-Src may play a role in events associated with both neuronal differentiation and maintenance of mature neuronal cell functions.

## CHROMOSOMAL LOCATION

Genetic locus: SRC (human) mapping to 20q11.23; Src (mouse) mapping to 2 H1.

## SOURCE

p-c-Src (H-8) is a mouse monoclonal antibody specific for an epitope containing Tyr 530 phosphorylated c-Src of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2b</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-166859 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## APPLICATIONS

p-c-Src (H-8) is recommended for detection of Tyr 530 phosphorylated c-Src of human and rat origin, and Tyr 535 phosphorylated c-Src of mouse origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for c-Src siRNA (h): sc-29228, c-Src siRNA (m): sc-29859, c-Src siRNA (r): sc-270199, c-Src shRNA Plasmid (h): sc-29228-SH, c-Src shRNA Plasmid (m): sc-29859-SH, c-Src shRNA Plasmid (r): sc-270199-SH, c-Src shRNA (h) Lentiviral Particles: sc-29228-V, c-Src shRNA (m) Lentiviral Particles: sc-29859-V and c-Src shRNA (r) Lentiviral Particles: sc-270199-V.

Molecular Weight of p-c-Src: 60 kDa.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent), Lambda Phosphatase: sc-200312A and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



Western blot analysis of c-Src phosphorylation in untreated (A,D), serum starved and EGF treated (B,E) and serum starved, EGF and lambda protein phosphatase treated (C,F) HEK293 whole cell lysates. Antibodies tested include p-c-Src (H-8): sc-166859 (A,B,C) and c-Src (17AT28): sc-130124 (D,E,F).



p-c-Src (H-8): sc-166859. Immunoperoxidase staining of formalin fixed, paraffin-embedded human testis tissue showing cytoplasmic and nuclear staining of cells in seminiferous ducts and cytoplasmic staining of Leydig cells (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human skin tissue showing cytoplasmic and nuclear staining of keratinocytes, fibroblasts, Langerhans cells and melanocytes (B).

## SELECT PRODUCT CITATIONS

- Choi, C.W., et al. 2015. Focal adhesion kinase and Src expression in premalignant and malignant skin lesions. *Exp. Dermatol.* 24: 361-364.
- Kung, M.L., et al. 2016. Nanoscale characterization illustrates the cisplatin-mediated biomechanical changes of B16-F10 melanoma cells. *Phys. Chem. Chem. Phys.* 18: 7124-7131.
- Lee, N.K., et al. 2024. Development of advanced cardiac progenitor cell culture system through fibronectin and vitronectin derived peptide coated plate. *Stem Cell Res.* 79: 103476.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.