



TRAIL (h2): 293T Lysate: sc-170196

BACKGROUND

Proteins belonging to the tumor necrosis factor (TNF) superfamily are potent mediators of the inflammation and immune systems. Members of the TNF superfamily include TNF β , lymphotoxin β (LT β), CD40L, CD30L, CD27L, Ox40L, 4-1BBL and FAS-L (APO-1). Most TNF family members are type II transmembrane proteins that are proteolytically processed at their carboxy-terminal extracellular domain to form a soluble homotrimeric molecule. The extracellular domain of an additional TNF family member, designated TNF-related apoptosis-inducing ligand (TRAIL) or APO-2L, exhibits 14-28% homology with other members of the TNF family. Like soluble FAS-L, soluble TRAIL will induce apoptosis. The morphological and cellular changes caused by the introduction of soluble TRAIL to Jurkat cells are indistinguishable from those caused by the introduction of soluble FAS-L. Unlike FAS-L, whose expression is more or less restricted to activated T cells, significant levels of TRAIL are observed in many tissues and it is constitutively expressed by some cell lines.

REFERENCES

1. Smith, C.A., Farrah, T. and Goodwin, R.G. 1994. The TNF receptor superfamily of cellular and viral proteins: activation, costimulation and death. *Cell* 76: 959-962.
2. Cosman, D. 1994. A family of ligands for the TNF receptor superfamily. *Stem Cells* 12: 440-455.
3. Cleveland, J.L. and Ihle, J.N. 1995. Contenders in FAS-L/TNF death signaling. *Cell* 81: 479-482.
4. Nagata, S. and Golstein, P. 1995. The FAS death factor. *Science* 267: 1449-1456.
5. Wiley, S.R., Schooley, K., Smolak, P.J., Din, W.S., Huang, C.P., Nicholl, J.K., Sutherland, G.R., Smith, T.D., Rauch, C. and Smith, C.A. 1995. Identification and characterization of a new member of the TNF family that induces apoptosis. *Immunity* 3: 673-682.
6. Baker, S.J. and Reddy, E.P. 1996. Transducers of life and death: TNF receptors superfamily and associated proteins. *Oncogene* 12: 1-9.
7. Pitti, R.M., Marsters, S.A., Ruppert, S., Donahue, C.J., Moore, A. and Ashkenazi, A. 1996. Induction of apoptosis by APO-2 ligand, a new member of the TNF cytokine family. *J. Biol. Chem.* 271: 12687-12690.
8. Plasilova, M., Zivny, J., Jelinek, J., Neuwirtova, R., Cermak, J., Necas, E., Andera, L. and Stopka, T. 2002. TRAIL (APO-2L) suppresses growth of primary human leukemia and myelodysplasia progenitors. *Leukemia* 16: 67-73.
9. Hyer, M.L., Croxton, R., Krajewska, M., Krajewski, S., Kress, C.L., Lu, M., Suh, N., Sporn, M.B., Cryns, V.L., Zapata, J.M. and Reed, J.C. 2005. Synthetic triterpenoids cooperate with TNF-related apoptosis-inducing ligand to induce apoptosis of breast cancer cells. *Cancer Res.* 65: 4799-4808.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: TNFSF10 (human) mapping to 3q26.31.

PRODUCT

TRAIL (h2): 293T Lysate represents a lysate of human TRAIL transfected 293 cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

TRAIL (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive TRAIL antibodies. Recommended use: 10-20 μ l per lane.

Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.