



GPI (h2): 293T Lysate: sc-170846

BACKGROUND

Glucose-6-phosphate isomerase (GPI) has many other names, including Phosphohexose isomerase (PHI), Neuroleukin (NLK) and Spermatigen-36 (SA-36). GPI is a cytoplasmic homodimer belonging to the GPI family. It is a neurotrophic factor for spinal and sensory neurons and is involved in glycolysis and gluconeogenesis. Defects or mutations in GPI can cause hereditary nonspherocytic hemolytic anemia (HA), hydrops fetalis, immediate neonatal death and neurological impairment.

REFERENCES

1. Beutler, E., West, C., Britton, H.A., Harris, J. and Forman, L. 1997. Glucosephosphate isomerase (GPI) deficiency mutations associated with hereditary nonspherocytic hemolytic anemia (HNSHA). *Blood Cells Mol. Dis.* 23: 402-409.
2. Kugler, W., Breme, K., Laspe, P., Muirhead, H., Davies, C., Winkler, H., Schroter, W. and Lakomek, M. 1998. Molecular basis of neurological dysfunction coupled with haemolytic anaemia in human glucose-6-phosphate isomerase (GPI) deficiency. *Hum. Genet.* 103:450-454.
3. Schulz, L.C. and Bahr, J.M. 2003. Glucose-6-phosphate isomerase is necessary for embryo implantation in the domestic ferret. *Proc. Natl. Acad. Sci. USA* 100: 8561-8566.
4. Muraki, Y., Matsumoto, I., Chino, Y., Hayashi, T., Suzuki, E., Goto D., Ito, S., Murata, H., Tsutsumi, A. and Sumida, T. 2004. Glucose-6-phosphate isomerase variants play a key role in the generation of anti-GPI antibodies: possible mechanism of autoantibody production. *Biochem. Biophys. Res. Commun.* 323: 518-522.
5. Graham Solomons, J.T., Zimmerly, E.M., Burns, S., Krishnamurthy, N., Swan, M.K., Krings, S., Muirhead, H., Chirgwin, J. and Davies, C. 2004. The crystal structure of mouse phosphoglucose isomerase at 1.6Å resolution and its complex with glucose 6-phosphate reveals the catalytic mechanism of sugar ring opening. *J. Mol. Biol.* 342: 847-860.
6. Schubert, D., Maier, B., Morawietz, L., Krenn, V. and Kamradt, T. 2004. Immunization with glucose-6-phosphate isomerase induces T cell-dependent peripheral polyarthritis in genetically unaltered mice. *J. Immunol.* 172: 4503-4509.
7. SWISS-PROT/TrEMBL (P06744). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

CHROMOSOMAL LOCATION

Genetic locus: GPI (human) mapping to 19q13.11.

PRODUCT

GPI (h2): 293T Lysate represents a lysate of human GPI transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

GPI (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive GPI antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.