

# Nrf1 (h2): 293T Lysate: sc-171305

## BACKGROUND

The NF-E2 DNA binding protein is composed of two subunits, p45 and MafK, and it regulates expression of globin genes in developing erythroid cells through interaction with Maf recognition elements (MAREs). A family of NF-E2 related proteins, which are collectively known as the Cap "n" collar (CNC) family and include Nrf1 (also designated TCF11), Nrf2 and Nrf3, are bZIP transcription factors that heterodimerize with Maf proteins to bind MARE sequences. The Nrf proteins also bind the antioxidant response element (ARE) and are implicated in the regulation of detoxification enzymes and the oxidative stress response. They do so by heterodimerizing with Jun family members (c-Jun, JunB and JunD) to activate gene expression, specifically the detoxifying enzyme, NQO1. Nrf2 is widely expressed and is thought to translocate to the nucleus after treatment with xenobiotics and antioxidants, which stimulate its release from a repressor protein Keap1. Nrf3 is highly expressed in placenta, B cells and monocytes.

## REFERENCES

1. Chan, J.Y., et al. 1995. Chromosomal localization of the human NF-E2 family of bZIP transcription factors by fluorescence *in situ* hybridization. *Hum. Genet.* 95: 265-269.
2. Chan, K., et al. 1996. Nrf2, a member of the NF-E2 family of transcription factors, is not essential for murine erythropoiesis, growth, and development. *Proc. Natl. Acad. Sci. USA* 93: 13943-13948.
3. Kobayashi, A., et al. 1999. Molecular cloning and functional characterization of a new cap "n" collar family transcription factor Nrf3. *J. Biol. Chem.* 274: 6443-6452.
4. Dhakshinamoorthy, S., et al. 2000. Small Maf (MafG and MafK) proteins negatively regulate antioxidant response element-mediated expression and antioxidant induction of the NAD(P)H:Quinone oxidoreductase1 gene. *J. Biol. Chem.* 275: 40134-40141.

## CHROMOSOMAL LOCATION

Genetic locus: NFE2L1 (human) mapping to 17q21.32.

## PRODUCT

Nrf1 (h2): 293T Lysate represents a lysate of human Nrf1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

Nrf1 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Nrf1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Nrf1 (E-4): sc-365651 is recommended as a positive control antibody for Western Blot analysis of enhanced human Nrf1 expression in Nrf1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



Nrf1 (E-4): sc-365651. Western blot analysis of Nrf1 expression in non-transfected: sc-117752 (A) and human Nrf1 transfected: sc-171305 (B) 293T whole cell lysates

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.