

CD1A (h2): 293T Lysate: sc-171320

BACKGROUND

The CD1 multigene family encodes five forms of the CD1 T cell surface glycoprotein in human, designated CD1A, 1B, 1C, 1D and 1E. CD1, a type 1 membrane protein, has structural similarity to the MHC class I antigen and has been shown to present lipid antigens for recognition by T lymphocytes. CD1 antigens are associated with β -2-Microglobulin and expressed on cortical thymocytes, Langerhans cells, a B cell subset and some dendritic cells. Specifically, CD1A is a marker for Langerhans cell histiocytosis (LCH) and is found on interdigitating cells. Adaptor-protein complexes and CD1-associated chaperones control CD1 trafficking and the development and activation of CD1-restricted T cells. Constitutive endocytosis of CD1B molecules and the differential sorting of MHC class II from lysosomes separate peptide- and lipid antigen-presenting molecules during dendritic cell maturation. CD1B is also expressed in interdigitating cells. The human CD1 genes are all closely linked in a cluster mapping at chromosome 1q23.1.

REFERENCES

1. Martin, L.H., Calabi, F., Lefebvre, F.A., Bilisland, C.A. and Milstein, C. 1987. Structure and expression of the human thymocyte antigens CD1A, CD1B, and CD1C. *Proc. Natl. Acad. Sci. USA* 84: 9189-9193.
2. Aruffo, A. and Seed, B. 1989. Expression of cDNA clones encoding the thymocyte antigens CD1A, B, C demonstrates a hierarchy of exclusion in fibroblasts. *J. Immunol.* 143: 1723-1730.
3. Longley, J., Kraus, J., Alonso, M. and Edelson, R. 1989. Molecular cloning of CD1A (T6), a human epidermal dendritic cell marker related to class I MHC molecules. *J. Invest. Dermatol.* 92: 628-631.
4. Sotzik, F., Boyd, A. and Shortman, K. 1993. Surface antigens of human thymocyte populations defined by CD3, CD4 and CD8 expression: CD1A is expressed by mature thymocytes but not peripheral T cells. *Immunol. Lett.* 36: 101-106.
5. Porcelli, S.A. 1995. The CD1 family: a third lineage of antigen-presenting molecules. *Adv. Immunol.* 59: 1-18.
6. Melian, A., Beckman, E.M., Porcelli, S.A. and Brenner, M.B. 1996. Antigen presentation by CD1 and MHC-encoded class I-like molecules. *Curr. Opin. Immunol.* 8: 82-88.
7. Storkus, W.J., Wei, M., Cresswell, P. and Dawson, J.R. 1996. Class I-like CD1A-C do not protect target cells from NK-mediated cytotoxicity. *Cell. Immunol.* 167: 154-156.
8. Bauer, A., Huttinger, R., Staffler, G., Hansmann, C., Schmidt, W., Majdic, O., Knapp, W. and Stockinger, H. 1997. Analysis of the requirement for β -2-Microglobulin for expression and formation of human CD1 antigens. *Eur. J. Immunol.* 27: 1366-1373.
9. Brigl, M. and Brenner, M.B. 2004. CD1: antigen presentation and T cell function. *Annu. Rev. Immunol.* 22: 817-890.

CHROMOSOMAL LOCATION

Genetic locus: CD1A (human) mapping to 1q23.1.

PRODUCT

CD1A (h2): 293T Lysate represents a lysate of human CD1A transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

CD1A (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive CD1A antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

CD1A (CTB6): sc-5265 is recommended as a positive control antibody for Western Blot analysis of enhanced human CD1A expression in CD1A transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



CD1A (CTB6): sc-5265. Western blot analysis of CD1A expression in non-transfected: sc-117752 (A) and human CD1A transfected: sc-171320 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.