



POR1 (h): 293T Lysate: sc-171384

BACKGROUND

POR1 (also designated arfapin 2) was first isolated as a Rac 1 binding protein necessary for Rac-mediated Actin polymerization and the subsequent formation of membrane ruffles and lamellipodia. POR1 has also been shown to interact with the ADP ribosylation factor ARF6, a GTPase that associates with the plasma membrane and intracellular endosome vesicles, in a GTP dependent manner. The association of POR1 with ARF6 stimulates induction of Actin polymerization. POR1 appears to play a regulatory role through multiple signaling pathways in the reorganization of the cytoskeletal structure.

REFERENCES

1. Van Aelst, L., Joneson, T. and Bar-Sagi, D. 1996. Identification of a novel Rac 1-interacting protein involved in membrane ruffling. *EMBO J.* 15: 3778-3786.
2. Joneson, T., McDonough, M., Bar-Sagi, D. and Van Aelst, L. 1996. Rac regulation of Actin polymerization and proliferation by a pathway distinct from Jun kinase. *Science* 274: 1374-1376.
3. D'Souza-Schorey, C., Boshans, R.L., McDonough, M., Stahl, P.D. and Van Aelst, L. 1997. A role for POR1, a Rac 1-interacting protein, in ARF6-mediated cytoskeletal rearrangements. *EMBO J.* 16: 5445-5454.
4. D'Souza-Schorey, C., van Donselaar, E., Hsu, V.W., Yang, C., Stahl, P.D. and Peters, P.J. 1998. ARF6 targets recycling vesicles to the plasma membrane: insights from an ultrastructural investigation. *J. Cell Biol.* 140: 603-616.
5. Gauthier-Rouviere, C., Vignal, E., Meriane, M., Roux, P., Montcourier, P. and Fort, P. 1998. Rho G GTPase controls a pathway that independently activates Rac 1 and Cdc42Hs. *Mol. Biol. Cell* 9: 1379-1394.
6. Radhakrishna, H., Al-Awar, O., Khachikian, Z. and Donaldson, J.G. 1999. ARF6 requirement for Rac ruffling suggests a role for membrane trafficking in cortical Actin rearrangements. *J. Cell Sci.* 112: 855-866.

CHROMOSOMAL LOCATION

Genetic locus: ARFIP2 (human) mapping to 11p15.4.

PRODUCT

POR1 (h): 293T Lysate represents a lysate of human POR1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

POR1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive POR1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.