# karyopherin $\alpha$ 1 (h3): 293 Lysate: sc-171440



The Boures to Overtion

#### **BACKGROUND**

Protein transport across the nucleus is a selective, multi-step process involving several cytoplasmic factors. Proteins must be recognized as import substrates, dock at the nuclear pore complex and translocate across the nuclear envelope in an ATP-dependent fashion. Two cytosolic factors centrally involved in the recognition and docking process are the karyopherin  $\alpha 1$  and karyopherin  $\beta 1$  subunits. Karyopherin  $\alpha 1$  functions in the recognition and targeting of substrates destined for nuclear import, while karyopherin  $\beta 1$  serves as an adapter, tethering the karyopherin  $\alpha 1/\text{substrate}$  complex to docking proteins on the nuclear envelope termed nucleoporins. Karyopherin  $\alpha 2$  has been shown to complex with Epstein-Barr virus nuclear antigen 1 (EBNA1). Certain RNA-binding proteins are imported to the nucleus by karyopherin  $\beta 2$ , and karyopherin  $\beta 3$  appears to be involved in the import of some ribosomal proteins.

# **REFERENCES**

- 1. Moroianu, J., et al. 1995. Previously identified protein of uncertain function is karyopherin  $\alpha$  and together with karyopherin  $\beta$  docks import substrate at nuclear pore complexes. Proc. Natl. Acad. Sci. USA 92: 2008-2011.
- Moroianu, J., et al. 1995. Protein export from the nucleus requires the GTPase Ran and GTP hydrolysis. Proc. Natl. Acad. Sci. USA 92: 4318-4322.
- 3. Lounsbury, K.M., et al. 1996. Ran binding domains promote the interaction of Ran with  $p97/\beta$ -karyopherin, linking the docking and translocation steps of nuclear import. J. Biol. Chem. 271: 2357-2360.
- 4. Moroianu, J., et al. 1996. The binding site of karyopherin  $\alpha$  for karyopherin  $\beta$  overlaps with a nuclear localization sequence. Proc. Natl. Acad. Sci. USA 93: 6572-6576.
- 5. Moroianu, J., et al. 1996. Nuclear protein import: Ran-GTP dissociates the karyopherin  $\alpha/\beta$  heterodimer by displacing  $\alpha$  from an overlapping binding site on  $\beta$ . Proc. Natl. Acad. Sci. USA 93: 7059-7062.
- 6. Fischer, N., et al. 1997. Epstein-Barr virus nuclear antigen 1 forms a complex with the nuclear transporter karyopherin  $\alpha$ 2. J. Biol. Chem. 272: 3999-4005.
- 7. Yaseen, N.R., et al. 1997. Cloning and characterization of human karyopherin β3. Proc. Natl. Acad. Sci. USA 94: 4451-4456.
- 8. Bonifaci, N., et al. 1997. Karyopherin  $\beta 2$  mediates nuclear import of a mRNA binding protein. Proc. Natl. Acad. Sci. USA 94: 5055-5060.

# CHROMOSOMAL LOCATION

Genetic locus: KPNA1 (human) mapping to 3q21.1.

# **PRODUCT**

karyopherin  $\alpha 1$  (h3): 293 Lysate represents a lysate of human karyopherin  $\alpha 1$  transfected 293 cells and is provided as 100  $\mu g$  protein in 200  $\mu l$  SDS-PAGE buffer.

# **STORAGE**

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

#### **APPLICATIONS**

karyopherin  $\alpha$ 1 (h3): 293 Lysate is suitable as a Western Blotting positive control for human reactive karyopherin  $\alpha$ 1 antibodies. Recommended use: 10-20  $\mu$ 1 per lane.


Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

karyopherin  $\alpha$ 1 (187.1): sc-101292 is recommended as a positive control antibody for Western Blot analysis of enhanced human karyopherin  $\alpha$ 1 expression in karyopherin  $\alpha$ 1 transfected 293 cells (starting dilution 1:100, dilution range 1:100-1:1,000).

#### **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG $\kappa$  BP-HRP: sc-516102 or m-lgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>TM</sup> Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

# **DATA**



karyopherin  $\alpha$ 1 (187.1): sc-101292. Western blot analysis of karyopherin  $\alpha$ 1 expression in nontransfected: sc-110760 ( $\bf A$ ) and human karyopherin  $\alpha$ 1 transfected: sc-171440 ( $\bf B$ ) 293 whole cell lysates.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com