

TCEAL1 (h): 293T Lysate: sc-171670

BACKGROUND

TCEAL1 (transcription elongation factor A (SII) protein 3), also known as p21, SIIR or pp21, is a member of the TFS-II family. Transcription elongation factors of the TFS-II family are responsible for releasing RNA polymerase II (Pol II) from transcriptional arrest. DNA arresting sites can result in locked ternary complexes if elongating RNA polymerases are trapped. Transcription elongation factors function to activate the intrinsic RNA cleavage activity of RNA polymerases. This allows the RNA polymerase to cleave the nascent transcript, thereby forming a new 3'-terminus to resume elongation. TCEAL1 is a 157 amino acid nuclear protein that is ubiquitously expressed. TCEAL1 may exert its effects via protein-protein interactions with other transcriptional regulators rather than via direct binding of DNA. Phosphorylation of TCEAL1 on Ser-36 and Ser-37 is critical for transcriptional repression.

REFERENCES

1. Yeh, C.H. and Shatkin, A.J. 1994. A HeLa-cell-encoded p21 is homologous to transcription elongation factor SII. *Gene* 143: 285-287.
2. Yeh, C.H. and Shatkin, A.J. 1994. Down-regulation of Rous sarcoma virus long terminal repeat promoter activity by a HeLa cell basic protein. *Proc. Natl. Acad. Sci. USA* 91: 11002-11006.
3. Gu, W. and Reines, D. 1995. Variation in the size of nascent RNA cleavage products as a function of transcript length and elongation competence. *J. Biol. Chem.* 270: 30441-30447.
4. Labhart, P. and Morgan, G.T. 1998. Identification of novel genes encoding transcription elongation factor TFIIS (TCEA) in vertebrates: conservation of three distinct TFIIS isoforms in frog, mouse, and human. *Genomics* 52: 278-288.
5. Pillutla, R.C., Shimamoto, A., Furuichi, Y. and Shatkin, A.J. 1999. Genomic structure and chromosomal localization of TCEAL1, a human gene encoding the nuclear phosphoprotein p21/SIIR. *Genomics* 56: 217-220.
6. Wind, M. and Reines, D. 2000. Transcription elongation factor SII. *Bioessays* 22: 327-336.
7. Kim, Y.B., Ki, S.W., Yoshida, M. and Horinouchi, S. 2000. Mechanism of cell cycle arrest caused by histone deacetylase inhibitors in human carcinoma cells. *J. Antibiot.* 53: 1191-1200.
8. Rodriguez, A., Jung, E.J., Yin, Q., Cayrol, C. and Flemington, E.K. 2001. Role of c-Myc regulation in Zta-mediated induction of the cyclin-dependent kinase inhibitors p21 and p27 and cell growth arrest. *Virology* 284: 159-169.
9. Makino, H., Tajiri, T., Miyashita, M., Sasajima, K., Anbazhagan, R., Johnston, J. and Gabrielson, E. 2005. Differential expression of TCEAL1 in esophageal cancers by custom cDNA microarray analysis. *Dis. Esophagus* 18: 37-40.

CHROMOSOMAL LOCATION

Genetic locus: TCEAL1 (human) mapping to Xq22.2.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

TCEAL1 (h): 293T Lysate represents a lysate of human TCEAL1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

TCEAL1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive TCEAL1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

TCEAL1 (F-4): sc-393621 is recommended as a positive control antibody for Western Blot analysis of enhanced human TCEAL1 expression in TCEAL1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



TCEAL1 (F-4): sc-393621. Western blot analysis of TCEAL1 expression in non-transfected: sc-117752 (A) and human TCEAL1 transfected: sc-171670 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.