



PICK1 (h4): 293T Lysate: sc-171856

BACKGROUND

Protein interacting with C kinase 1 (PICK1) is a PDZ-domain containing protein that is located in the perinuclear region and is phosphorylated in response to PKC α activation. PKC α , which is essential for the regulation of proliferation and differentiation in numerous cell types, contains within its catalytic region a PDZ-binding domain that is absent from other PKC isoforms. Mutation of the PICK1 PDZ domain inhibits the binding of PICK1 to PKC α . PICK1 also interacts with the carboxy-terminus of α -amino-3-hydroxy-5-methyl-isoxazole-4-propionic acid (AMPA) receptor, a neurotransmitter receptor located at excitatory synapses, suggesting that PICK1 plays a role in the modulation of synaptic transmission by targeting and anchoring AMPA to specific synapses.

REFERENCES

1. Staudinger, J., Zhou, J., Burgess, R., Elledge, S.J. and Olson, E.N. 1995. PICK1: a perinuclear binding protein and substrate for protein kinase C isolated by the yeast two-hybrid system. *J. Cell Biol.* 128: 263-271.
2. Staudinger, J., Lu, J. and Olson, E.N. 1997. Specific interaction of the PDZ domain protein PICK1 with the COOH terminus of protein kinase C- α . *J. Biol. Chem.* 272: 32019-32024.
3. Xia, J., Zhang, X., Staudinger, J. and Huganir, R.L. 1999. Clustering of AMPA receptors by the synaptic PDZ domain-containing protein PICK1. *Neuron* 22: 179-187.
4. Wyszynski, M., Valtschanoff, J.G., Naisbitt, S., Dunah, A.W., Kim, E., Standaert, D.G., Weinberg, R. and Sheng, M. 1999. Association of AMPA receptors with a subset of glutamate receptor-interacting protein *in vivo*. *J. Neurosci.* 19: 6528-6537.
5. Dev, K.K., Nishimune, A., Henley, J.M. and Nakanishi, S. 1999. The protein kinase C α binding protein PICK1 interacts with short but not long form alternative splice variants of AMPA receptor subunits. *Neuropharmacology* 38: 635-644.
6. Matsuzawa, D., Hashimoto, K., Miyatake, R., Shirayama, Y., Shimizu, E., Maeda, K., Suzuki, Y., Mashimo, Y., Sekine, Y., Inada, T., Ozaki, N., Iwata, N., Harano, M., Komiyama, T., Yamada, M., Sora, I., Ujike, H., et al. 2007. Identification of functional polymorphisms in the promoter region of the human PICK1 gene and their association with methamphetamine psychosis. *Am. J. Psychiatry* 164: 1105-1114.
7. Pan, L., Wu, H., Shen, C., Shi, Y., Jin, W., Xia, J. and Zhang, M. 2007. Clustering and synaptic targeting of PICK1 requires direct interaction between the PDZ domain and lipid membranes. *EMBO J.* 26: 4576-4587.
8. Zhang, C.S., Bertaso, F., Eulenburg, V., Lerner-Natoli, M., Herin, G.A., Bauer, L., Bockaert, J., Fagni, L., Betz, H. and Scheschonka, A. 2008. Knock-in mice lacking the PDZ-ligand motif of mGluR7a show impaired PKC-dependent autoinhibition of glutamate release, spatial working memory deficits, and increased susceptibility to pentylenetetrazol. *J. Neurosci.* 28: 8604-8614.
9. Haglerod, C., Kapic, A., Boulland, J.L., Hussain, S., Holen, T., Skare, O., Laake, P., Ottersen, O.P., Haug, F.M. and Davanger, S. 2009. Protein interacting with C kinase 1 (PICK1) and GluR2 are associated with presynaptic plasma membrane and vesicles in hippocampal excitatory synapses. *Neuroscience* 158: 242-522.

CHROMOSOMAL LOCATION

Genetic locus: PICK1 (human) mapping to 22q13.1.

PRODUCT

PICK1 (h4): 293T Lysate represents a lysate of human PICK1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

PICK1 (h4): 293T Lysate is suitable as a Western Blotting positive control for human reactive PICK1 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.