



Rfp2 (h2): 293T Lysate: sc-172517

BACKGROUND

The tripartite motif (TRIM) family of proteins are characterized by a conserved TRIM domain that includes a coiled-coil region, a B box-type zinc finger, one RING finger and three zinc-binding domains. Rfp2 (Ret finger protein 2), also known as TRIM13 (tripartite motif-containing 13), CAR, RNF77 or LEU5, is a 407 amino acid protein that belongs to the TRIM protein family and contains one B box-type zinc finger and one RING-type zinc finger. Existing as two alternatively spliced isoforms designated α and β , Rfp2 is thought to act as a tumor suppressor that, when defective, may be involved in the development and progression of B-cell chronic lymphocytic leukemia. Additionally, Rfp2 may function as an E3 ubiquitin ligase that is involved in protein degradation pathways related to the ER-associated degradation (ERAD) pathway.

REFERENCES

1. Liu, Y., et al. 1993. Chronic lymphocytic leukemia cells with allelic deletions at 13q14 commonly have one intact RB1 gene: evidence for a role of an adjacent locus. *Proc. Natl. Acad. Sci. USA* 90: 8697-8701.
2. Liu, Y., et al. 1995. 13q deletions in lymphoid malignancies. *Blood* 86: 1911-1915.
3. Kapanadze, B., et al. 1998. A cosmid and cDNA fine physical map of a human chromosome 13q14 region frequently lost in B-cell chronic lymphocytic leukemia and identification of a new putative tumor suppressor gene, Leu5. *FEBS Lett.* 426: 266-270.
4. Raymond, A., et al. 2001. The tripartite motif family identifies cell compartments. *EMBO J.* 20: 2140-2151.
5. van Everdink, W.J., et al. 2003. Rfp2, c13ORF1, and FAM10A4 are the most likely tumor suppressor gene candidates for B-cell chronic lymphocytic leukemia. *Cancer Genet. Cytogenet.* 146: 48-57.
6. Corcoran, M.M., et al. 2004. DLEU2 encodes an antisense RNA for the putative bicistronic Rfp2/LEU5 gene in humans and mouse. *Genes Chromosomes Cancer* 40: 285-297.
7. Skoblov, M., et al. 2006. Human Rfp2 gene promoter: unique structure and unusual strength. *Biochem. Biophys. Res. Commun.* 342: 859-866.
8. Lerner, M., et al. 2007. The RBCC gene RFP2 (Leu5) encodes a novel transmembrane E3 ubiquitin ligase involved in ERAD. *Mol. Biol. Cell* 18: 1670-1682.
9. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 605661. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: TRIM13 (human) mapping to 13q14.2.

PRODUCT

Rfp2 (h2): 293T Lysate represents a lysate of human Rfp2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Rfp2 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Rfp2 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.