

eIF2B β (h2): 293T Lysate: sc-172832

BACKGROUND

The initiation of protein synthesis in eukaryotic cells is regulated by interactions between protein initiation factors and RNA molecules. The eukaryotic initiation complex eIF2B exists as a five subunit complex composed of eIF2 α , eIF2 β , eIF2 γ , eIF2B δ , and eIF2B ϵ . The eIF2B complex catalyzes the exchange of GDP for GTP on the eIF2 complex, following the interaction of eIF2/GTP with the 40S ribosomal subunit. Guanine nucleotide exchange factor (GEF) activity is exhibited by the eIF2B ϵ subunit alone, but is greater in the presence of all five eIF2B subunits. Phosphorylation of eIF2 inhibits GEF activity of eIF2B, an inhibition that requires the eIF2 α subunit.

REFERENCES

- Trachsel, H. and Staehelin, T. 1978. Binding and release of eukaryotic initiation factor eIF2 and GTP during protein synthesis initiation. *Proc. Natl. Acad. Sci. USA* 75: 204-208.
- Benne, R., Ames, H., Hershey, J.W. and Voorma, H.O. 1979. The activity of eukaryotic initiation factor eIF2 in ternary complex formation with GTP and Met-tRNA. *J. Biol. Chem.* 254: 3201-3205.
- Ernst, H., Duncan, R.F. and Hershey, J.W. 1987. Cloning and sequencing of complementary DNAs encoding the α -subunit of translational initiation factor eIF2. Characterization of the protein and its messenger RNA. *J. Biol. Chem.* 262: 1206-1212.
- Pathak, V.K., Nielsen, P.J., Trachsel, H. and Hershey, J.W. 1988. Structure of the β subunit of translational initiation factor eIF2. *Cell* 54: 633-639.
- Kaufman, R.J., Davies, M.V., Pathak, V.K. and Hershey, J.W. 1989. The phosphorylation state of eucaryotic initiation factor 2 alters translational efficiency of specific mRNAs. *Mol. Cell. Biol.* 9: 946-958.
- Gaspar, N.J., Kinzy, T.G., Scherer, B.J., Humbelin, M., Hershey, J.W. and Merrick, W.C. 1994. Translation initiation factor eIF2. Cloning and expression of the human cDNA encoding the γ subunit. *J. Biol. Chem.* 269: 3415-3422.

CHROMOSOMAL LOCATION

Genetic locus: EIF2B2 (human) mapping to 14q24.3.

PRODUCT

eIF2B β (h2): 293T Lysate represents a lysate of human eIF2B β transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

eIF2B β (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive eIF2B β antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

eIF2B β (E-12): sc-376478 is recommended as a positive control antibody for Western Blot analysis of enhanced human eIF2B β expression in eIF2B β transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



eIF2B β (E-12): sc-376478. Western blot analysis of eIF2B β expression in non-transfected: sc-117752 (A) and human eIF2B β transfected: sc-172832 (B) 293T whole cell lysates.



eIF2B β (P-4): sc-9979. Western blot analysis of eIF2B β expression in non-transfected: sc-117752 (A) and human eIF2B β transfected: sc-172832 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.