

USP3 (h3): 293T Lysate: sc-173209

BACKGROUND

The ubiquitin (Ub) pathway involves three sequential enzymatic steps that facilitate the conjugation of Ub and Ub-like molecules to specific protein substrates. Through the use of a wide range of enzymes that can add or remove ubiquitin, the Ub pathway controls many intracellular processes such as signal transduction, transcriptional activation and cell cycle progression. USP3 (ubiquitin specific peptidase 3), also known as UBP or SIH003, is a 520 amino acid protein that contains one UBP-type zinc finger and belongs to the peptidase C19 family. Expressed ubiquitously with highest levels present in pancreas, USP3 catalyzes the conversion of a ubiquitin C-terminal thioester to a free ubiquitin and a thiol. The gene encoding USP3 maps to human chromosome 15, which houses over 700 genes and comprises nearly 3% of the human genome.

REFERENCES

1. Chung, C.H. and Baek, S.H. 1999. Deubiquitinating enzymes: their diversity and emerging roles. *Biochem. Biophys. Res. Commun.* 266: 633-640.
2. Sloper-Mould, K.E., Eyre, H.J., Wang, X.W., Sutherland, G.R. and Baker, R.T. 1999. Characterization and chromosomal localization of USP3, a novel human ubiquitin-specific protease. *J. Biol. Chem.* 274: 26878-26884.
3. Puente, X.S., Sánchez, L.M., Overall, C.M. and López-Otín, C. 2003. Human and mouse proteases: a comparative genomic approach. *Nat. Rev. Genet.* 4: 544-558.
4. Online Mendelian Inheritance in Man, OMIM[™]. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 604728. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Whalen, R., Liu, X. and Boyer, T.D. 2006. Identification of a short form of ubiquitin-specific protease 3 that is a repressor of rat glutathione S-transferase gene expression. *Biochem. J.* 394: 519-526.
6. Nicassio, F., Corrado, N., Vissers, J.H., Areces, L.B., Bergink, S., Marteijn, J.A., Geverts, B., Houtsmuller, A.B., Vermeulen, W., Di Fiore, P.P. and Citterio, E. 2007. Human USP3 is a chromatin modifier required for S phase progression and genome stability. *Curr. Biol.* 17: 1972-1977.

CHROMOSOMAL LOCATION

Genetic locus: USP3 (human) mapping to 15q22.31.

PRODUCT

USP3 (h3): 293T Lysate represents a lysate of human USP3 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

USP3 (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive USP3 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

USP3 (8L8): sc-135597 is recommended as a positive control antibody for Western Blot analysis of enhanced human USP3 expression in USP3 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



USP3 (8L8): sc-135597. Western blot analysis of USP3 expression in non-transfected: sc-117752 (A) and human USP3 transfected: sc-173209 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.