



FcRH3 (h3): 293T Lysate: sc-173229

BACKGROUND

The Fc receptor homolog (FcRH) family of proteins are related to the classical Fc receptors (FcR) and belong to the immunoglobulin receptor superfamily. The proteins in the FcRH family are type I transmembrane glycoproteins and consist of FcRH1-FcRH6. The genes encoding the human FcRH proteins map to chromosome 1, near the related FCR genes. The FcRH proteins, which are involved in immune system regulation, have immunoreceptor-tyrosine inhibitory motifs in their cytoplasmic domains. Mutations in the gene encoding for FcRH proteins may be associated with systemic lupus erythematosus, autoimmune thyroid disease and rheumatoid arthritis. The highly restricted pattern of mouse FcRH3 expression suggests this member of the phylogenetically conserved FcRH family may have an important immunoregulatory role in marginal zone B cells.

REFERENCES

1. Davis, R.S., Wang, Y.H., Kubagawa, H. and Cooper, M.D. 2001. Identification of a family of Fc receptor homologs with preferential B cell expression. *Proc. Natl. Acad. Sci. USA* 98: 9772-9777.
2. Davis, R.S., Dennis, G., Jr., Odom, M.R., Gibson, A.W., Kimberly, R.P., Burrows, P.D. and Cooper, M.D. 2002. Fc receptor homologs: newest members of a remarkably diverse Fc receptor gene family. *Immunol. Rev.* 190: 123-136.
3. Ehrhardt, G.R., Davis, R.S., Hsu, J.T., Leu, C.M., Ehrhardt, A. and Cooper, M.D. 2003. The inhibitory potential of Fc receptor homolog 4 on memory B cells. *Proc. Natl. Acad. Sci. USA* 100: 13489-13494.
4. Davis, R.S., Stephan, R.P., Chen, C.C., Dennis, G., Jr. and Cooper, M.D. 2004. Differential B cell expression of mouse Fc receptor homologs. *Int. Immunol.* 16: 1343-1353.
5. Davis, R.S., Ehrhardt, G.R., Leu, C.M., Hirano, M. and Cooper, M.D. 2005. An extended family of Fc receptor relatives. *Eur. J. Immunol.* 35: 674-680.

CHROMOSOMAL LOCATION

Genetic locus: FCRL3 (human) mapping to 1q23.1.

PRODUCT

FcRH3 (h3): 293T Lysate represents a lysate of human FcRH3 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

FcRH3 (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive FcRH3 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.